



Facultative adjustment of the offspring sex ratio and male attractiveness: a systematic review and meta-analysis

Booksmythe, Isobel ; Mautz, Brian ; Davis, Jacqueline ; Nakagawa, Shinichi ; Jennions, Michael D

Abstract: Females can benefit from mate choice for male traits (e.g. sexual ornaments or body condition) that reliably signal the effect that mating will have on mean offspring fitness. These male-derived benefits can be due to material and/or genetic effects. The latter include an increase in the attractiveness, hence likely mating success, of sons. Females can potentially enhance any sex-biased benefits of mating with certain males by adjusting the offspring sex ratio depending on their mate's phenotype. One hypothesis is that females should produce mainly sons when mating with more attractive or higher quality males. Here we perform a meta-analysis of the empirical literature that has accumulated to test this hypothesis. The mean effect size was small ($r = 0.064$ – 0.095 ; i.e. explaining $<1\%$ of variation in offspring sex ratios) but statistically significant in the predicted direction. It was, however, not robust to correction for an apparent publication bias towards significantly positive results. We also examined the strength of the relationship using different indices of male attractiveness/quality that have been invoked by researchers (ornaments, behavioural displays, female preference scores, body condition, male age, body size, and whether a male is a within-pair or extra-pair mate). Only ornamentation and body size significantly predicted the proportion of sons produced. We obtained similar results regardless of whether we ran a standard random-effects meta-analysis, or a multi-level, Bayesian model that included a correction for phylogenetic non-independence. A moderate proportion of the variance in effect sizes (51.6–56.2%) was due to variation that was not attributable to sampling error (i.e. sample size). Much of this non-sampling error variance was not attributable to phylogenetic effects or high repeatability of effect sizes among species. It was approximately equally attributable to differences (occurring for unknown reasons) in effect sizes among and within studies (25.3, 22.9% of the total variance). There were no significant effects of year of publication or two aspects of study design (experimental/observational or field/laboratory) on reported effect sizes. We discuss various practical reasons and theoretical arguments as to why small effect sizes should be expected, and why there might be relatively high variation among studies. Currently, there are no species where replicated, experimental studies show that mothers adjust the offspring sex ratio in response to a generally preferred male phenotype. Ultimately, we need more experimental studies that test directly whether females produce more sons when mated to relatively more attractive males, and that provide the requisite evidence that their sons have higher mean fitness than their daughters.

DOI: <https://doi.org/10.1111/brv.12220>

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-124247>
Journal Article

Originally published at:

Booksmythe, Isobel; Mautz, Brian; Davis, Jacqueline; Nakagawa, Shinichi; Jennions, Michael D (2017). Facultative adjustment of the offspring sex ratio and male attractiveness: a systematic review and meta-analysis. *Biological Reviews of the Cambridge Philosophical Society*, 92(1):108-134.
DOI: <https://doi.org/10.1111/brv.12220>

3

4 **Facultative adjustment of the offspring sex ratio and male**
5 **attractiveness: a systematic review and meta-analysis**

6

7 Isobel Booksmythe^{1,2,*}, Brian Mautz^{1,2}, Jacqueline Davis^{1,3}, Shinichi Nakagawa^{4,5} and
8 Michael D. Jennions¹

9

10 ¹*Research School of Biology, The Australian National University, Canberra, ACT 0200, Australia*

11 ²*Department of Animal Ecology, Evolutionary Biology Centre, Uppsala University, Norbyvägen*
12 *18D, SE-75236 Uppsala, Sweden*

13 ³*Department of Psychology, University of Cambridge, Downing Street, Cambridge, CB2 3EB, U.K.*

14 ⁴*Department of Zoology, University of Otago, PO Box 56, Dunedin, New Zealand*

15 ⁵*Evolution & Ecology Research Centre and School of Biological, Earth and Environmental Sciences,*
16 *University of New South Wales, Sydney, NSW 2052, Australia*

17

18 *Author for correspondence (E-mail: isobel.booksmythe@ieu.uzh.ch; Tel.: +41 044 6354804).

19

20 **ABSTRACT**

21 Females can benefit from mate choice for male traits (e.g. sexual ornaments or body condition) that
22 reliably signal the effect that mating will have on mean offspring fitness. These male-derived benefits
23 can be due to material and/or genetic effects. The latter include an increase in the attractiveness,
24 hence likely mating success, of sons. Females can potentially enhance any sex-biased benefits of
25 mating with certain males by adjusting the offspring sex ratio depending on their mate’s phenotype.
26 One hypothesis is that females should produce mainly sons when mating with more attractive or
27 higher quality males. Here we perform a meta-analysis of the empirical literature that has

28 accumulated to test this hypothesis. The mean effect size was small ($r = 0.064\text{--}0.095$; i.e. explaining
29 $<1\%$ of variation in offspring sex ratios) but statistically significant in the predicted direction. It was,
30 however, not robust to correction for an apparent publication bias towards significantly positive
31 results. We also examined the strength of the relationship using different indices of male
32 attractiveness/quality that have been invoked by researchers (ornaments, behavioural displays, female
33 preference scores, body condition, male age, body size, and whether a male is a within-pair or extra-
34 pair mate). Only ornamentation and body size significantly predicted the proportion of sons
35 produced. We obtained similar results regardless of whether we ran a standard random-effects meta-
36 analysis, or a multi-level, Bayesian model that included a correction for phylogenetic non-
37 independence. A moderate proportion of the variance in effect sizes ($51.6\text{--}56.2\%$) was due to
38 variation that was not attributable to sampling error (i.e. sample size). Much of this non-sampling
39 error variance was not attributable to phylogenetic effects or high repeatability of effect sizes among
40 species. It was approximately equally attributable to differences (occurring for unknown reasons) in
41 effect sizes among and within studies (25.3% , 22.9% of the total variance). There were no significant
42 effects of year of publication or two aspects of study design (experimental/observational or
43 field/laboratory) on reported effect sizes. We discuss various practical reasons and theoretical
44 arguments as to why small effect sizes should be expected, and why there might be relatively high
45 variation among studies. Currently, there are no species where replicated, experimental studies show
46 that mothers adjust the offspring sex ratio in response to a generally preferred male phenotype.
47 Ultimately, we need more experimental studies that test directly whether females produce more sons
48 when mated to relatively more attractive males, *and* that provide the requisite evidence that their sons
49 have higher mean fitness than their daughters.

50

51 *Key words:* female choice, mate choice, maternal effects, sex allocation, sex ratio adjustment, sexual
52 selection.

53

54 CONTENTS

55 I. Introduction

56 (1) Environmental condition-dependent sex allocation

57 (2) Genetic condition-dependent sex ratio adjustment

58 (3) Models of sex ratio adjustment (SRA) in response to mate attractiveness

59 (4) The empirical evidence

60 (5) The role of meta-analysis

61 II. Methods

62 (1) Literature search

63 (2) Data extraction and coding

64 (3) Standard random-effects meta-analyses

65 (4) Multilevel meta-analyses including phylogenetic effects

66 (5) Bias detection

67 III. Results

68 (1) Attrition of studies

69 (2) Standard random-effects meta-analyses

70 (3) Multilevel meta-analyses including phylogenetic effects

71 IV. Discussion

72 (1) General findings

73 (2) Reasons for a small effect: sex allocation decisions are influenced by many factors

74 (3) Reasons for a small effect: measuring sexual selection is tricky

75 (4) Updated results and future improvements

76 V. Conclusions

77 VI. Acknowledgements

78 VII. References

79 VIII. Supporting information

80

81 **I. INTRODUCTION**

82 **(1) Environmental condition-dependent sex allocation**

83 The birth sex ratio in many species deviates from unity, despite selection on parents to produce equal
84 numbers of sons and daughters due to negative frequency-dependent selection (Düsing, 1884; Fisher,
85 1930). Although the reproductive value of individuals of the rare sex is always greater than that of
86 the other sex, thereby tending to select for equal investment into each sex, this does not constrain the

87 offspring sex ratio to be even. First, if one sex is cheaper to produce and total sex allocation is equal,
88 more offspring of the cheaper sex will be produced (West, 2009). Second, adaptive adjustment of sex
89 allocation away from equal investment into both sexes is favoured by selection if the function
90 relating investment to fitness returns differs for sons and daughters (i.e. if it is non-linear for at least
91 one sex: Charnov, 1979) even if there is no variation in the resources mothers can allocate to
92 reproduction (reviews: Frank, 1990, 1998 p.174–178). Third, facultative parental control of the
93 offspring sex ratio is potentially advantageous when offspring fitness depends on some aspect of the
94 breeding environment, which varies among parents (e.g. territory quality, female condition, adult
95 size), if this condition dependence differently affects the absolute reproductive value of sons and
96 daughters (Trivers & Willard, 1973). This is likely, as many environmental factors have sex-specific
97 effects on offspring fitness (e.g. Jones, Nakagawa & Sheldon, 2009; Bowers, Thompson & Sakaluk,
98 2014). More generally, it is clear that males and females tend to have different fitness functions
99 (review: Connallon & Clark, 2014). Given variation among parents in the quality of their breeding
100 environment, those breeding in a relatively better environment should increase allocation towards the
101 sex that has greater reproductive value when investment is higher. One way to do this is to bias the
102 offspring sex ratio towards that sex.

103 Trivers & Willard (1973) originally suggested that differences in maternal condition could lead to
104 facultative sex ratio adjustment (henceforth, SRA) by parents. They noted that in polygynous
105 mammals (e.g. many ungulates and pinnipeds) there is enormous variation in male mating success,
106 and that larger, healthier males are disproportionately successful at acquiring mates when males
107 compete. By contrast, female size has weak effects on fecundity (in many mammals the range in litter
108 size is very low). In general, mothers in good physical condition have the ability to produce larger,
109 healthier offspring. Trivers & Willard (1973) therefore argued that mothers in good condition should
110 preferentially produce sons because of the relatively greater fitness gains they accrue from improving
111 their sons' body condition. There is empirical support for this claim from studies of ungulates (meta-
112 analysis: Sheldon & West, 2004).

Hypotheses of condition-dependent SRA have since been broadened to include many other environmental and social factors that affect ‘breeding conditions’ and generate predictable differences in the relative fitness of sons and daughters (West, 2009, chapter 6). These scenarios include: production of daughters by high-ranking mothers when social rank is maternally inherited, as in primates (Schino, 2004); production of daughters by parasitoid insects laying eggs on larger hosts because of the stronger effect of size on female fecundity than on male mating success (meta-analysis: West & Sheldon, 2002); production of the helping sex if parents lack sufficient helpers-at-the-nest in cooperatively breeding birds and mammals (meta-analysis: West & Sheldon, 2002; but see Stubblefield & Orzack, 2013); seasonal or laying order effects if one sex requires more resources as a neonate, and food availability changes seasonally (e.g. for work on raptors see Rutz, 2012 and references therein). More generally, many factors have been posited to affect the optimal offspring sex ratio because they predict future resource availability for investment into offspring (e.g. territory quality, harem status; for a more complete list see West, 2009, p.183).

(2) Genetic condition-dependent sex ratio adjustment

There is an important condition-dependent sex allocation scenario that is not based on parental ability to predict environmentally determined resource availability. It is the argument that a male’s attractiveness or quality can differentially affect the fitness of his sons and daughters because of genetic effects (Burley, 1981, 1986). Males with larger ornaments and faster display rates, those in better body condition and/or those of larger body size are often preferred by females, gaining above-average mating success (Andersson, 1994). There is strong evidence for moderate heritability of sexually selected, preferred male traits (Prokuda & Roff, 2014; Wyman & Rowe, 2014). Burley (1981, 1986) therefore predicted that females mating with attractive males should preferentially produce sons because sons will benefit more than daughters by inheriting genes that increase their sexual attractiveness. Burley’s idea is compelling because few traits are more intimately related to reproductive success and, by extension, to fitness, than those under directional sexual selection

(Shuster & Wade, 2003). There are also plausible grounds for assuming that some of the benefits of mating with attractive males are sex-specific, or sex-biased, accruing only or mainly to sons. Still, there are several important caveats.

Preferred male traits can signal both material and genetic benefits (e.g. Hoelzer, 1989; meta-analysis: Møller & Jennions, 2001). If there are direct, fecundity-enhancing benefits of female choice due to material gains, then additional maternal resources might benefit daughters as much as, or even more than, sons (e.g. if the fecundity gains from increasing female size outweigh those of higher mating success for larger males or males who can invest more into costly ornaments). The adaptive prediction that females will produce sons when mated to attractive males implicitly assumes there are indirectly selected, genetic benefits of mate choice. Furthermore, for genetic benefits to favour SRA towards sons, the genes inherited from attractive sires must disproportionately advantage males (Burley, 1986). This is intuitively the case for arbitrarily attractive (Fisherian) male traits that are not inherited by daughters. The inheritance of genes that increase viability (so-called ‘good genes’) must, however, have a stronger effect on males than females to favour a male-biased offspring sex ratio. This might occur, for example, if sons benefit more than daughters from improved body condition because many sexual ornaments have highly condition-dependent expression (Cotton, Fowler & Pomiankowski, 2004). The benefits of an attractive father can also be greater for sons than daughters if attractiveness is similarly heritable for sons and daughters (i.e. both sexes are under sexual selection due to mate choice), but males experience greater variance in reproductive success (e.g. Du & Lu, 2010; Bowers *et al.*, 2013), which is causally related to mate choice (i.e. sexual selection is stronger on males than females). In general, selection will favour a conditional sex allocation strategy when ornamentation imparts heritable benefits (be they due to the ornaments themselves or traits genetically correlated with ornamentation) that are greater for offspring of one sex than for the other (Fawcett *et al.*, 2011).

If male-biased genetic benefits of mate choice exist, they can be enhanced by biasing the offspring sex ratio towards sons when mating with more attractive males, and towards daughters when mating

with less attractive males (Burley, 1981, 1986). This form of conditional SRA has several potential fitness benefits for mothers that are driven by genetic effects. (1) It increases the production of ‘sexy sons’ (Weatherhead & Robertson, 1979; Firman, 2011; Prokop *et al.*, 2012). There is, however, an implicit assumption that the benefits of a higher mating rate are not perfectly countered by greater naturally selected costs (e.g. higher mortality) (see Pen & Weissing, 2000), or countered by a trade-off with other sexually selected traits (e.g. reduced sperm competitiveness; see Lüpold *et al.*, 2014). (2) Producing fewer daughters reduces the negative effects of sexually antagonistic genes when mated to an attractive male (if genes from more attractive sires make daughters of below-average fitness) (Rice & Chippindale, 2001; review: Rice, 2013). (3) Even in the absence of any such sexual antagonism, producing more daughters lessens the disadvantage of mating with an unattractive male as the value of daughters increases due to their ‘underproduction’ by females mated to attractive mates. Consequently, the fitness difference between females mated to attractive and unattractive mates declines (Fawcett *et al.*, 2007, 2011).

(3) Models of sex ratio adjustment (SRA) in response to mate attractiveness

Greater production of sons due to genes that confer male-specific fitness benefits is distinct from most other conditional sex allocation scenarios. In the latter, the factors that favour biased sex ratios (e.g. maternal condition, habitat quality) tend to do so because they predict the availability of resources that can be invested into offspring, and the sexes differ in the effect of investment on their reproductive value (Frank, 1998). Consequently, most theoretical treatments of conditional sex allocation are not readily applicable to the mate attractiveness hypothesis. This is largely because covariances between genes for different traits (e.g. ornaments, preferences and sex allocation strategies), and changes in genetic variances have to be taken into account when modelling genetic benefits. Crucially, selection on condition-dependent SRA does not obviously affect variation in parental resource availability that favours SRA, and the sex-specific functions relating reproductive value to environmental quality remain the same. By contrast, selection on SRA in response to

additive genetic variation in attractiveness could lead to a co-evolutionary feedback that changes the fitness benefits of being more attractive because it will affect phenotypic variation in male attractiveness and female mating preferences (Fawcett *et al.*, 2011).

To date, only four mathematical models have addressed the evolution of SRA in response to male attractiveness. Pen & Weissing (2000) provided the first formal model. They used a game theory approach with a discrete male trait (i.e. ornamented *versus* unornamented males) to show that an evolutionarily stable strategy for male-biased SRA when mated to ornamented males could evolve. The stability of such a strategy was, however, sensitive to the form of selection driving male trait exaggeration. Facultative SRA evolved under ‘good genes’ selection where the male trait signalled heritable viability (albeit in a simplified scenario where ‘viability’ genes did not benefit daughters). It did not evolve under a pure Fisherian process. In the latter case, all males are of equal fitness at equilibrium as natural selection perfectly balances sexual selection (ornamented males have greater mating success but higher mortality). There was therefore no fitness benefit from SRA towards sons when mated to an attractive male in this model.

Subsequently Fawcett *et al.* (2007) ran individual-based simulations that confirmed these initial results. Facultative SRA did not evolve when the male ornamental trait was purely Fisherian, but did evolve when it was a conditional indicator of viability, and when it was a Fisherian trait subject to a mutation bias (*sensu* Pomiankowski, Iwasa & Nee, 1991). Reassuringly, the results did not differ between two different proximate mechanisms of SRA. The findings were, however, slightly different if the male trait exhibited continuous rather than discrete variation. For a continuous male trait, the only scenario where facultative SRA unambiguously evolved was for a Fisherian trait subject to mutation bias. Although there was still a shift in sex allocation when the male trait was a conditional indicator, it was weak and inconsistent across simulation runs. The evolution of a stronger sex ratio bias for a Fisherian trait makes intuitive sense because such traits only benefit males. By contrast, conditional indicators signal the presence of viability genes that increase the fitness of both sexes. An additional, clear finding of Fawcett *et al.* (2007) was that the evolution of SRA in response to male

217 attractiveness was in all cases very slow, suggesting that it is under very weak selection. This is
218 readily seen by comparing the rate of evolution of facultative SRA to the rate of change in the female
219 preference and expression of preferred male traits in the simulations (Fawcett *et al.*, 2007). Selection
220 in the wild is likely to be even weaker than reported in the models, because (a) the models ignore the
221 additional costs of actually adjusting the sex ratio (e.g. energetic costs if this requires strategic
222 reabsorption of embryos, or time costs of delaying fertilization; Krackow, 1995); and (b) in practice,
223 females obtain imperfect information on male genotype due to both non-heritable environmental
224 influences on male sexual trait expression (Holman & Kokko, 2013), and female perceptual errors
225 when assessing the expression of sexual signals (e.g. Farris & Ryan, 2011).

226

227 More recently, Fawcett *et al.* (2011) noted that facultative SRA in response to heritable sire
228 attractiveness reduces sexual selection on male traits. A co-evolutionary feedback occurs because
229 females mated to less attractive males ‘compensate’ by producing daughters. This reduces the fitness
230 difference between females mated to more and less attractive males. The strength of selection for
231 costly female preferences for males with more elaborate sexual traits is therefore weakened. In
232 addition, sexual selection declines because the mean strength of the female preference in the
233 population is lower when choosier females (those mating with more attractive males) produce fewer
234 daughters, because only daughters express mating preferences. Both processes select for reduced
235 investment into costly sexual traits by males. Facultative SRA effectively reduces the heritability of
236 attractiveness and mating preferences because of the lower fitness gain of ornamentation (Fawcett *et*
237 *al.*, 2011). Fawcett *et al.* (2011) therefore conclude that species with elaborate sexual traits are the
238 least likely to show facultative SRA in response to mate attractiveness. This is worrisome because
239 empirical researchers testing for adaptive SRA have sought out species with strong sexual selection
240 on males.

241 Booksmythe, Schwanz & Kokko (2013) suggested that the feedback between sexual selection and
242 SRA based on attractiveness might be even more complicated than indicated by Fawcett *et al.* (2011).

Specifically, as the expression of a preferred male trait declines, it is increasingly difficult for females to distinguish and discriminate between males due to perceptual errors while mate sampling. More generally, imperfect maternal control of sex allocation is expected. They also noted that given costs of adjusting the sex ratio, this strategy might be eliminated when male trait expression is low. In an extension of Fawcett *et al.*'s (2011) model they allowed the 'accuracy' of sex allocation to evolve (i.e. the extent to which there is a 'step-like' transition between the production of daughters and sons; see West, Reece & Sheldon, 2002, for comments on empirical evidence that sex allocation is imprecise and not 'step-like'). They also imposed a cost on females adjusting the sex ratio that increased with the precision of sex allocation. The simulation results were complex, suggesting there is not a simple linear relationship between the strength of SRA and male trait expression. In general, the greatest degree of SRA arose when sexual selection on male traits was intermediate in strength.

254

255 **(4) The empirical evidence**

Many empirical studies testing Burley's original idea have accumulated in the last three decades (Fig. 1), but the available evidence for adjustment of the offspring sex ratio in response to mate attractiveness is still mixed. An early meta-analysis found a strong positive correlation between the proportion of sons in a brood and male attractiveness based on 11 avian studies of eight species ($r = 0.205$; West & Sheldon, 2002). This result seemed to spur a flurry of further research, but the extent to which the hypothesis is currently supported is less clear. Several studies have found a significant positive relationship between some measure of male attractiveness and the proportion of sons in a brood (e.g. Saino *et al.*, 2002; Sato & Karino, 2010; Correa *et al.*, 2011; Bowers *et al.*, 2013; others have found no such relationship (Ramsay *et al.*, 2003; Rutstein *et al.*, 2004; Parker, 2005; Limbourg, Mateman & Lessells, 2013), and still others have reported a significant relationship in the opposite direction (Rutstein *et al.*, 2005; Delhey *et al.*, 2007). In addition, studies over multiple years or from different populations of the same species have reported that the relationship varies in strength and/or direction (e.g. Griffith *et al.*, 2003; Rosivall *et al.*, 2004; see also Parker, 2013, for a meta-analysis of

studies on blue tits). The extent to which intra-specific variation is due to sampling error or genuine biological differences is unknown.

Ultimately, the diverse empirical results obtained to date reflect the challenge of identifying male traits that: (a) have a sound theoretical reason to affect the offspring sex ratio (rather than other forms of differential allocation such as preferential provisioning of eggs or juveniles of one sex; see Horvátová, Nakagawa & Uller, 2012), (b) are associated with male-biased genetic benefits, and (c) are applicable to the study population/species. In combination with the added difficulties of controlling for confounding variables, dealing with measurement error, and recurrent problems of low statistical power in behavioural ecology (Jennions & Møller, 2003) it is encouraging that any studies have obtained a significant result in the predicted direction. Even so, determining whether this has occurred more often than expected by chance remains a major challenge.

280

281 **(5) The role of meta-analysis**

Qualitative, narrative reviews have reported mixed evidence for adaptive sex allocation in birds and mammals, regardless of the source of selection on the offspring sex ratio (reviews: Clutton-Brock, 1986; Sheldon, 1998; Palmer, 2000; West, Herre & Sheldon 2000; West *et al.*, 2002; Komdeur & Pen, 2002). By contrast, formal meta-analyses of hypotheses related to sex allocation in birds, mammals and insects have produced more clear-cut conclusions. Several meta-analyses have reported that the average relationship is significant in the predicted direction [e.g. helpers-at-the-nest: West & Sheldon, 2002; Griffin, Sheldon & West, 2005; but see Stubblefield & Orzack, 2013; mate attractiveness in birds: West & Sheldon, 2002; but see Parker, 2013; maternal condition in ungulates: Sheldon & West, 2004; host size in parasitoids: West & Sheldon, 2002; mate quality, laying date and season in birds: Cassey, Ewen & Møller, 2006 (correction of Ewen, Cassey & Møller, 2004); local mate competition: West, Shuker & Sheldon, 2005; split sex ratios in social insects: Meunier, West & Chapuisat, 2008; local resource competition in primates: Silk & Brown, 2008]. However, other meta-analyses, often using similar data sets, have reported that the mean relationship (effect size) is not

295 significantly different from the null expectation (zero) (e.g. maternal rank in primates: Brown & Silk,
296 2002; Schino, 2004; Silk, Willoughby & Brown, 2005; maternal condition in mammals: Cameron,
297 2004). Meta-analysis is a statistical technique designed to detect broad-scale patterns across empirical
298 studies and quantitatively weigh studies by their accuracy in reporting the focal relationship. In short,
299 estimates of a relationship are made with clearly stated confidence intervals. By contrast, narrative
300 reviews are susceptible to vote counting (how many studies obtained a ‘yes’ or ‘no’ answer), which
301 ignores the limitations of low power that plague most studies in evolutionary ecology. Meta-analysis
302 has the additional advantage that it provides tools to gauge the extent to which moderating factors are
303 associated with the observed relationships, and tools to assess the likely influence of publication
304 biases (Jennions *et al.*, 2012). For example, although the mean effect size did not differ from zero in
305 Ewen *et al.* (2004), moderator variables had a significant influence on offspring sex ratios (Cassey *et*
306 *al.*, 2006). This suggests that, under certain conditions, there is still evidence that parents can skew
307 sex ratios.

308 As new empirical data accumulate, meta-analyses need to be revised. This is especially important in
309 fields where effect sizes tend to decline over time (see Jennions & Møller, 2002). For example,
310 Pereira & Ioannidis (2011) have recently estimated – based on a decline in the reported mean effect
311 sizes between meta-analyses conducted in 2005 and 2010 – that 37% of medical meta-analyses are
312 likely to conclude falsely that there is evidence for a relationship. In addition, it is obvious to anyone
313 who has conducted a meta-analysis that subjective decisions are made (e.g. how to extract effect
314 sizes, how to search for papers). This can lead to researchers reporting different findings from meta-
315 analyses on the same question, even when they use similar data sets. For a good example, see a recent
316 dispute about the effect of the menstrual cycle on human mate choice between Wood *et al.* (2014)
317 and Gildersleeve, Haselton & Fales (2014). For a topic closer to the current issue one can compare
318 the findings of two meta-analyses on SRA in relation to helper number by Griffin *et al.* (2005) and
319 Stubblefield & Orzack (2013).

320 The most recent meta-analysis of Burley's (1981) hypothesis was by West & Sheldon (2002). Since
321 then, there have been many studies with data on SRA in relation to mate attractiveness/quality (Fig.
322 1). The accumulation of studies allows for a more robust test of whether there is a general trend,
323 which seems especially relevant in the light of recent theory questioning the extent to which there
324 will be detectable selection for SRA based on male attractiveness (Fawcett *et al.*, 2011). Here we use
325 both standard meta-analytical and newer meta-regression approaches (including a control for
326 phylogenetic relationships) to: (1) test whether there is a general trend for the offspring sex ratio to be
327 male-biased when females are mated to attractive males. (2) Investigate if the relationship between
328 the offspring sex ratio and male attractiveness depends on the type of male trait or measurement used
329 as a proxy for attractiveness. For example, are ornaments better predictors of SRA than behaviour?
330 (3) Test whether phylogenetic effects and differences in study design are associated with the strength
331 of the relationship between the offspring sex ratio and male attractiveness.

332

333 **II. METHODS**

334 **(1) Literature search**

335 We used two complementary approaches to identify as many relevant studies as possible. First, we
336 performed a key word search using the *ISI Web of Knowledge* search engine (for details of the
337 databases covered see online supporting information, Appendix S1). We cross-referenced a set of key
338 words related to sex ratio (e.g. "offspring sex ratio", "proportion male") to two sets of key words
339 relating to male attractiveness: (1) general terms (e.g. attractive, ornament, "secondary sexual trait")
340 and (2) numerous specific secondary sexual characteristics (e.g. plumage, antler, horn, badge). The
341 exact search term combinations used are given in Appendix S1. Second, we identified six empirical
342 and theoretical papers that are highly cited in studies of sex allocation in response to mate
343 attractiveness: Burley (1981), Ellegren, Gustafsson & Sheldon (1996), Frank (1990), Fawcett *et al.*
344 (2007), Sheldon *et al.* (1999) and West & Sheldon (2002). At least one of these papers is likely to be
345 cited by a study testing for SRA in response to male attractiveness. We performed a forward search

to obtain papers citing any of these works. In total, the two search approaches yielded 4474 unique references (online Appendix S2). To finalize the data-checking and analysis protocols we conducted our final search on 17 July 2014. We did not add papers encountered by other means because these are more likely to be drawn from higher impact journals that might preferentially publish studies with significant results (Murtaugh, 2002). We also did not solicit unpublished data sets from colleagues. This procedure reduces the risk of biasing our estimates of effect sizes (see Jennions *et al.*, 2013). As with an empirical study, our goal was to obtain an extensive, representative sample of studies rather than to locate every available data point. The criteria we used to decide whether a study contained potentially usable data are provided in online Appendix S1. In brief, we excluded studies of plants, humans and farmed animals. We looked for studies where there were likely to be data on both brood/litter offspring sex ratios (i.e. not just population-level ratios) and measurements of male traits that are generally assumed to be targets of female choice in many taxa (e.g. body condition, body size and, most importantly, male ornamentation).

360

361 **(2) Data extraction and coding**

We collected and transformed study results into a standard effect size measure (Z_r ; see below) to allow for comparisons between studies. Reported effects were used in the following order of preference: (1) direct reports of an effect size (e.g. correlations) ($N = 13$); (2) effect measures with magnitude and direction (e.g. regression coefficients and standard errors, standardized mean differences, or descriptive statistics) ($N = 79$); (3) raw numbers (e.g. contingency tables) ($N = 12$); (4) inferential (test) statistics (often from multiple regressions) (e.g. t , z , F and χ^2 values, P values) ($N = 161$). In some cases we could only note that there was a non-significant effect ($N = 11$). All data were converted to correlation coefficients r , and then to Z_r , following the formulae listed in Lipsey & Wilson (2001) and Nakagawa & Cuthill (2007). The variance of Z_r is $1/(N-3)$ (Lipsey & Wilson, 2001). We calculated separate effect sizes for each measure of attractiveness from every independent

372 population and/or experiment (i.e. each unique group of parental individuals; hereafter referred to as
373 a ‘study’). Thus, if a paper tested the relationship between a male ornament and offspring sex ratio in
374 two distinct populations, these were considered independent studies and the paper contributed two
375 effect sizes to the meta-analysis. In some cases, where data were analysed by year, each year could
376 reasonably be considered an independent sample of individuals and was treated as a separate study
377 (e.g. Griffith *et al.*, 2003).

378 The included studies defined and measured male attractiveness in different ways. Namely, (a)
379 researchers chose focal traits based on evidence from previous research that particular morphological
380 or behavioural traits influenced male success in attracting and/or mating with females; (b) researchers
381 assumed that sexually dimorphic traits reflect sexual selection on males such that the trait will affect
382 male reproductive success; (c) researchers based their hypothesis on the general prediction that extra-
383 pair males are more attractive than average; (d) in a few studies researchers directly measured female
384 preferences (e.g. association time with different males) to score or rank test males. We were broad in
385 our criteria for what made an ‘attractive’ trait acceptable for inclusion as some studies cited strong
386 evidence for the effect on male mating success of a certain trait (e.g. experimental manipulation of
387 the trait affected female mate choice), while others would include this trait precisely because it had
388 been shown to influence sex ratios in other species. To test whether relationships varied across
389 different attractiveness proxies, including broad classes of male phenotypic traits, we grouped them
390 into seven categories: Ornament, Behaviour, Size, Condition, Preference, Age, and Extra-pair
391 paternity (EPP) (see Table 1). Each effect size was assigned to one of these categories. ‘Ornament’
392 included measures of the size or degree of elaboration of secondary sexual traits; for example, colour
393 measures or the size of a particular body part such as a comb, badge or elongated tail feather. Studies
394 that had manipulated male morphology (e.g. by adding coloured leg bands, or lengthening or
395 shortening tail feathers) were also included in this category. Similarly, ‘Behaviour’ included any
396 measures of male sexual behaviour that had been identified in the studies as attractive to females,
397 such as song or call attributes, courtship displays, and nest decoration collected by males. ‘Size’

398 included measures of structural body size and body mass. ‘Condition’ included measures of moult
 399 timing, parasite infection, and the residuals of mass on body size. ‘Preference’ measured
 400 attractiveness using female preference scores for particular males (e.g. association time in choice
 401 tests). Measures of male age or breeding experience were grouped into the category ‘Age’. Finally,
 402 EPP studies compared the sex ratio of within-pair and extra-pair offspring either within broods
 403 (extra-pair young *versus* within-pair young), between broods (i.e. broods with *versus* without EPP),
 404 or across broods (extra-pair and within-pair young pooled across all broods) (see Table 1).
 405 Studies could include multiple effect size estimates due to measurement of several male traits from
 406 the same category (e.g. two different ornaments) and/or from different categories (e.g. an ornament
 407 and a measure of condition). The presence of more than one effect size per study presents potential
 408 problems of non-independence of data for analyses. We dealt with this issue by running the meta-
 409 analysis in two ways. First, we used standard random-effects meta-analytic models, which require
 410 one effect-size estimate per study (Borenstein *et al.*, 2009). Here we calculated a mean effect size for
 411 each study by conducting a within-study meta-analysis (i.e. weighting the estimates by the sample
 412 size, as this occasionally differed even within a study). This was done separately for analyses
 413 restricted to a single trait type, or by pooling effect sizes across all trait types for analyses of ‘All’
 414 traits. Second, we ran multi-level, random-effects meta-analytic models, which allowed for the
 415 inclusion of multiple, non-independent effect size estimates per study by including the levels study
 416 and species (Nakagawa & Santos, 2012; Mengersen, Jennions & Schmid, 2013). In these analyses
 417 individual effect sizes were used and we included the trait type as a moderator.
 418 The effect size calculated was the Fisher-transformed correlation coefficient, Z_r . By convention,
 419 researchers refer to $r = 0.1$, 0.3 and 0.5 as small, medium and large effect sizes (Cohen, 1988). We
 420 present Z_r in our results, which is almost identical to r at low values (e.g. 0.1003 , 0.310 and 0.549 are
 421 the equivalent Z_r benchmarks). The effect size was coded as positive if an increase in the male
 422 attractiveness measure (i.e. greater ornamentation, more courtship, larger size, better condition,
 423 stronger preference score) was associated with a greater proportion of male offspring. We assumed

that older males were more attractive as this is a general assumption in bird studies (Møller & Ninni, 1998; Brooks & Kemp, 2001), and that extra-pair males were more attractive than within-pair males (Møller & Birkhead, 1994; but see Dunn, Whittingham & Pitcher, 2001; reviews: Westneat & Stewart, 2003; Forstmeier *et al.*, 2014). The variance in Z_r was calculated based on the number of broods in the study. We used the number of broods rather than unique breeding pairs as the sample size because this was the information most widely available across studies. Using different sources of N among studies to weight them is problematic because it gives greater weighting to studies that used lower-level analyses (e.g. an analysis that uses N = chicks). Where necessary, however, effect sizes were calculated using the source of N that was appropriate for the inferential statistic the authors presented. Notably, when we converted t or z values from mixed-effect models, we used N = broods rather than N = chicks (which potentially inflates the effect size; *cf.* Nakagawa & Cuthill, 2007; see also Section IV). We also coded information about the study that might have influenced the effect size, for use in a moderator analysis: study species, year of publication (z -transformed for analysis), study design (experimental or observational), and study location (laboratory or field research). There was substantial variation among studies in the stage at which offspring sex ratio was measured; we therefore included as a moderator ‘primary’ (measured at birth/laying) *versus* ‘secondary’ (some juvenile mortality had occurred) sex ratio rather than excluding studies on this basis. Seventeen papers reported insufficient information to calculate an effect size for one or more measured male traits (most commonly only reporting that the effect was ‘not significant’). We obtained the relevant information directly from the authors of 10 papers, retaining 11 unknown effects from the remaining seven papers. In addition, 28 papers contained one or more effects for which only model fit statistics or P values were reported, allowing us to extract the magnitude of an effect but not its direction. We obtained the direction of the effect from the authors of three of these papers. We retained 56 ‘directionless’ effects from the remaining 25 papers. We attempted to contact all authors so ‘incomplete’ data reflects no response by an author or their inability to recalculate the effect size (i.e. data files unavailable).

450 In all analyses, we tried to account for the uncertainty introduced by effects of unknown magnitude
451 and/or direction. First, we excluded all unknown and directionless effect sizes (dataset 1). Second, if
452 this analysis showed a significant mean effect, we added the unknown effect sizes and assigned them
453 a value of zero (i.e. assumed that they were highly ‘non-significant’) (dataset 2). Third, if the analysis
454 was still significant, we added the directionless effect sizes and also assigned them a value of zero
455 (dataset 3). Finally, if the analysis was still significant, we assigned all directionless effect sizes a
456 negative direction (dataset 4). Each successive dataset represents increasingly conservative
457 assumptions about the magnitude/direction of missing data. This allowed us to test the robustness of
458 our findings to the value of unknown and directionless effects.

459

460 **(3) Standard random-effects meta-analyses**

461 We conducted random-effects meta-analysis for all trait categories combined (the ‘All’ analysis), as
462 well as for each of the seven trait categories separately. These analyses assumed independence of
463 effect sizes. Each study therefore contributed one effect size per trait category (and thus per meta-
464 analysis). As noted above, if studies provided more than one effect size estimate per category, the
465 weighted mean effect for the category was calculated. Likewise, for the ‘All’ analysis we calculated
466 the weighted mean effect across all effect sizes for the study. We then assigned the variance to this
467 combined effect size based on the study sample size (number of broods). This is the most
468 conservative option as it assumes complete dependence between multiple estimates (e.g. that $r = 1$ for
469 the correlation between within-study estimates: see p. 228 in Borenstein *et al.*, 2009).

470 For all random-effects meta-analytic models, we used a restricted maximum likelihood (REML)
471 method to estimate τ^2 (the true between-study variance), which was combined with the within-study
472 variance for each effect size (Borenstein *et al.*, 2009). For the ‘All’ data set, we used meta-regression
473 to determine the effects of four study-level moderator variables: year of publication, design
474 (experimental/observational), location (field/laboratory), and stage at which sex ratio was measured

(primary/secondary) on the effect size. Based on the results, we did not include these moderators in the subsequent analyses for separate trait categories.

We also tested whether there was more heterogeneity in effect sizes among studies than could be explained by sampling error alone using Cochran's Q statistic. This formally tests whether there is greater variation in effect sizes among studies than expected if the true effect is identical for all studies (Hedges & Olkin, 1985). We also summarised heterogeneity using I^2 , which is often defined as the ratio of true heterogeneity to the total variance among studies. Borenstein *et al.* (2009) note, however, that this is not always correct because the ratio assumes a constant within-study variance, which is not the case (sampling error varies due to studies differing in sample size). They suggest that I^2 be treated as a measure of 'inconsistency' in effect sizes among studies. Higgins *et al.* (2003) have suggested that benchmarks for I^2 of 25%, 50% and 75% indicate low, moderate and high values of inconsistency among studies.

All statistical analyses for the above models were conducted in *R* statistical software using the *metafor* package (Viechtbauer, 2010).

(4) Multilevel meta-analyses including phylogenetic effects

While the standard meta-analyses described above included studies of vertebrate and invertebrate taxa, the multilevel meta-analyses with and without phylogenetic effects were limited to bird species, because there were few studies on other taxa in our data set (84.2% of species, 87.1% of included papers and 92.0% of effect sizes were for birds). The other taxa also had relatively few species sampled within each family. A tree including all species would be unreliable in terms of branch lengths (i.e. distance relationships between species). By contrast, we now have avian super-trees that include all extant species (Jezt *et al.*, 2012; see below). For birds, our data set had enough variation to conduct a phylogenetic meta-analysis. Studies could contribute multiple effect sizes because the multilevel model partially corrects for non-independence by modelling multiple random effects for both study and species. A complete correction of non-independence requires modelling the

501 correlations among effect sizes. For example, effect sizes for ornament and body size are likely to be
502 correlated when these traits were measured from the same group of individuals. Given the difficulty
503 of determining actual correlations among related effect sizes (as such correlations are almost always
504 not reported), the multilevel models we report in the main text assume correlations of zero.
505 Additionally, as supplementary materials (Table S2) we report on multilevel models that
506 conservatively assume all correlations to be 0.5. Note that qualitatively the results are very similar
507 among these models. As for the standard random-effect meta-analyses we initially excluded all
508 directionless and unknown effect sizes (avian dataset 1) and then added them to subsequent analyses
509 if the preceding analysis yielded a significant mean effect size (avian datasets 2–4).
510 To estimate the overall effect size we used four Bayesian Markov Chain Monte Carlo (MCMC)
511 generalised linear mixed-effects models, implemented in the MCMCglmm package of *R* (Hadfield,
512 2010; Hadfield & Nakagawa, 2010). First, the multilevel meta-analytic model (model A) included
513 study and species as random effects. We did not include ‘paper’ as a random effect because it closely
514 overlaps with study (99 and 81 levels, respectively, as most papers comprised a single study). This
515 model was analogous to the standard random-effects meta-analysis described above, but used MCMC
516 estimator rather than REML estimator and had two extra random effects (study and species). Second,
517 we extended model A by running a phylogenetic meta-analysis that included a phylogenetic tree
518 (model B). Third, we ran a multilevel meta-regression (model C) that extended model A by including
519 moderators (fixed effects): year of publication, seven categories of attractiveness measure (Ornament,
520 Behaviour, Size, Condition, Preference, Age, and EPP), design (experimental or observational), and
521 location (laboratory or field study). This was to assess whether these study characteristics impacted
522 the effect sizes, similar to the meta-regression used in the standard random-effects meta-analysis.
523 Finally, for completeness, we ran a meta-regression model (model D) that extended model C by
524 including phylogenetic effects. Models C and D were both run without the intercept. This ensures
525 that the mean estimates for the seven trait categories (conditional on the other fixed effects) were
526 directly obtained (for details, see Section III).

527 For both phylogenetic models (B and D) we used two different avian phylogenetic trees: (i) a version
 528 of the Ericson backbone trees from Jetz *et al.* (2012), referred to as the Ericson tree, and (ii) a version
 529 of the Hackett backbone trees also from Jetz *et al.* (2012), referred to as the Hackett tree. These
 530 phylogenies are presented as online supporting information in Fig. S1. Both trees were ultrametric
 531 with branch lengths. Each phylogenetic model produced quantitatively similar and qualitatively
 532 identical results using either tree. We therefore only present results for the Ericson tree (for results
 533 from the Hackett tree see Table S1 and Fig. S2). As an index of phylogenetic signal we calculated
 534 phylogenetic heritability H^2 , which can be defined as the proportion of phylogenetic variance in
 535 relation to the sum of all other variance components except sample error variance. Along with
 536 $I^2_{\text{phylogeny}}$ (see below), H^2 provides the magnitude of phylogenetic signal in the data. H^2 is equivalent
 537 to Pagel's λ when the unit of analysis is at the species level (Housworth, Martins & Lynch, 2004).
 538 For all multilevel (phylogenetic) models, we used an inverse Gamma prior ($V = 0.002$ and $nu = 1$) for
 539 all random effects. Each model was run for 1,100,000 iterations with a thinning interval of 1000 after
 540 a burn-in of 100,000 iterations. These settings resulted in posterior distributions consisting of 1000
 541 samples for all parameters. We assessed chain mixing by examining autocorrelation among posterior
 542 samples; they were less than 0.1 for all estimated parameters (Hadfield, 2010). We report our point
 543 estimates from models based on posterior means rather than posterior modes. We considered fixed
 544 factor estimates statistically significant if the 95% credible interval (CI, also known as the highest
 545 posterior density, which can be considered a Bayesian equivalent of the frequentist confidence
 546 interval) did not overlap zero. To quantify heterogeneity in multilevel meta-analytic models, we
 547 calculated a modified version of the I^2 statistics, following Nakagawa & Santos (2012). Basically,
 548 this partitions the proportion of 'unknown' variance not attributable to sampling variance (i.e.
 549 traditional I^2) into the contribution of random factors. In the current case these are the variance in
 550 effect sizes due to phylogenetic relatedness, differences among species, differences among studies,
 551 and differences in within-study variation (also called 'residual' variation). The sum of the

percentages of total variation due to these four sources equals the traditional I^2 of Higgins *et al.* (2003).

(5) Bias detection

A funnel plot reports the relationship between effect sizes and a measure of their variance (e.g. sampling variance, standard error or sample size). Variability in the observed effect size should decrease as sample size increases (i.e. as the estimate becomes more accurate due to lower sampling error). This yields a funnel-shaped plot. Almost all statistical tests for publication bias are based on the assumption that the funnel plot should be symmetrical (Jennions *et al.*, 2013). Asymmetry is assumed to reflect publication bias. We used funnel plots for visual inspection of funnel asymmetry, and then conducted Egger’s regression (Egger *et al.*, 1997) to test statistically for publication bias (i.e. funnel plot asymmetry) for each category of attractiveness measure. We then conducted trim-and-fill tests (the L0 model; Duval & Tweedie, 2000a, b) as another test for publication bias. Asymmetry due to publication bias is only predicted to appear as missing studies on the left of the plot (i.e. fewer significant results in the direction opposite to a true non-zero effect), but trim-and-fill can detect asymmetry on either side. We present results when studies were missing on the left or right. We do so because, in a broader context, the proportion of tests that identify missing studies on the left *versus* right can be viewed as an indicator of the extent to which asymmetry is due to publication bias (left only) rather than other factors that might generate asymmetry on either side. Small sample sizes limit the detection of significant asymmetry using Egger’s regression (Sterne & Egger, 2005), and funnel asymmetry can occur by chance especially when the sample size is low. Consequently, care is required when interpreting these publication bias analyses. Tests for bias were only run on ‘Data 1’ data sets (i.e. excluding unknown and directionless effect-size estimates). For the multi-level models (i.e. the avian-only data sets) we conducted analyses of bias on meta-analytic residuals (*sensu* Nakagawa & Santos, 2012). This ensures that we meet assumptions of independence, and that we reduce the effect of heterogeneity due to moderators that

could induce funnel asymmetry. All publication bias analyses were conducted using functions in the R package *metafor*.

III. RESULTS

(1) Attrition of studies

We scanned the title and abstract of all 4474 papers generated by our search protocol to identify studies potentially suitable for inclusion in the meta-analysis. We then read and assessed the methods and results sections of short-listed studies to determine their eligibility for inclusion. We identified 118 eligible papers. After full-text screening, 25 of these papers were excluded (Fig. 2, online Table S2). In total, 276 effect sizes for the meta-analysis were computed from 113 independent studies within 93 published papers. Of these, 254 effect sizes were from studies of birds (48 species). The remaining studies contributed nine effect sizes from four insect species, eight effect sizes from one species of fish, four effect sizes from three reptile species, and one effect size from a mammal. The full data set, and the data set reduced to one effect size per trait category per study are provided in online Appendix S3.

(2) Standard random-effects meta-analyses

The mean effect of male attractiveness on offspring sex ratio for traits pooled across all categories was small, positive and statistically significant for ‘dataset 1’ (Fig. 3A, Table 2). Females mated to attractive males produced more sons. This result was robust to the implementation of the successively more conservative datasets 2, 3 and 4 (all $P < 0.005$; Table 2). There were, however, reasons for concern about the effect of publication bias. The Egger’s regression test revealed significant asymmetry in the funnel plot ($t_{79} = 3.45$, $P = 0.001$; Table 3), and trim-and-fill estimated 17 (± 5.9 S.E.) studies ‘missing’ from the left-hand side of the plot (see Fig. 4A). Correcting for this bias substantially reduced the mean effect size estimate, which no longer differed from zero ($P = 0.149$; Table 3).

604 The heterogeneity due to variation between studies that could not be attributed to sampling error, I^2 ,
 605 was moderate at 56.20% (*sensu* Higgins *et al.*, 2003). Given the wide range of taxa, traits and study
 606 designs, one might expect a higher level of heterogeneity. To try to explain some of this
 607 heterogeneity, we ran a meta-regression model including the moderators year of publication, design,
 608 location and stage at which sex ratio was measured. They did not, however, explain a significant
 609 proportion of the observed heterogeneity ($Q_4 = 2.848$, $P = 0.584$; $b_{\text{year}} = 0.014$, 95% confidence
 610 interval, CI $[-0.324, 0.061]$, $b_{\text{Experiment-Observation}} = 0.026$ $[-0.080, 0.133]$, $b_{\text{Field-Laboratory}} = 0.082$ $[-$
 611 $0.022, 0.185]$, $b_{\text{Primary-Secondary}} = 0.014$ $[-0.082, 0.109]$). Given this result, we did not run meta-
 612 regression models for the other trait categories.

613 When we examined the effects for each trait category separately, we found a significantly positive,
 614 albeit small, effect for both male ornamentation and male size (Fig. 3A, Table 2). Females mated to
 615 more ornamented males or to larger males produced a higher proportion of sons than those mating
 616 with less ornamented or smaller mates. For both categories, this result held across datasets 1–3, and
 617 was only statistically non-significant in the most conservative ‘dataset 4’ analysis (where
 618 directionless effect sizes were assigned negative values) (Table 2). For ornaments, correction for a
 619 potential publication bias identified using trim-and-fill resulted in a mean effect size estimate that
 620 was non-significant ($P = 0.350$; Table 3). By contrast, for male size, trim-and-fill identified 7 (± 3.6
 621 S.E.) studies ‘missing’ from the right of the funnel plot. Correcting for this asymmetry increased the
 622 mean effect size estimate to 0.15 (± 0.03 S.E.; $P < 0.001$). The mean effect size estimate did not differ
 623 from zero for the other five trait types (behaviour, body condition, female preference score, male age,
 624 and male extra-pair or within-pair status) (Fig. 3A, Table 2). For male age, the Egger’s test suggested
 625 a publication bias, and trim-and-fill analysis suggested there were four missing studies. Even so, the
 626 addition of putative effect sizes still did not yield a significant effect size (Table 3). Trim-and-fill
 627 analyses also suggested there were three missing studies on the left for body condition, and two on
 628 the right for preference scores. Again, the addition of these studies did not yield significant mean
 629 effect size estimates (both $P > 0.089$; Table 3). Funnel plots are shown in Fig. 4.

The estimates of heterogeneity for the individual trait categories (dataset 1 analyses) ranged from $I^2 = 42.56$ to 73.43% (Table 2). This is moderate to high (*sensu* Higgins *et al.*, 2003), but it should be noted that uncertainty in estimates of I^2 is high when sample sizes are small. These results, along with those for all trait categories combined, suggest that over half the variation in effect sizes was due to systematic between-study differences. The sources of this variation are explored in Section IV. Notably, however, there was no heterogeneity in the effect size estimates for extra-pair mating status ($I^2 = 0$), indicating all variation was due to sampling error, although the number of studies in this analysis was rather limited.

(3) Multilevel meta-analyses including phylogenetic effects

For both the simple and phylogenetic multilevel meta-analyses we found small but significantly positive mean effect sizes for ‘avian dataset 1’ analyses (Table 4; Fig. 3A,B,C). These estimates were very similar to those from the standard random-effects meta-analysis for all traits pooled. The simple multilevel meta-analytic model (model A) had 95% credible intervals that did not span zero for avian datasets 2 and 3, but did for the most conservative dataset 4 (but even then, the lower CI was only just below zero at -0.002 ; Table 4). For the phylogenetic multilevel meta-analysis (model B), the mean effect had a 95% credible interval that included zero for avian dataset 2, so we did not run datasets 3 or 4.

In both multilevel meta-analytic models we can partition the variance not due to sampling error (i.e. heterogeneity). Here, I^2_{study} is the variance due to differences among studies while I^2_{species} is that among species. In the phylogenetic models, I^2_{species} is the variation due to the non-phylogenetic portion of species-level differences (associated with repeatability of effect sizes among species), while $I^2_{\text{phylogeny}}$ is the variance due to phylogenetic relatedness. $I^2_{\text{effect size}}$ is the residual variance that is not due to the sampling error. I^2_{total} is the sum of these values. In the simple multilevel meta-analysis, the largest component of variance for ‘avian dataset 1’ (effects of known direction and magnitude) was at the effect-size level (24.5%) then the study level (21.0%). Species had little

influence (6.0%) (Table 4), suggesting that repeatability is very low, although low replication within species (only 13 of 57 species were represented in more than one study, and only six species in more than two studies) likely contributes to this result. For the phylogenetic, multilevel meta-analysis of avian dataset 1, the largest component was also at the effect-size level (23.2%) followed by the study level (16.3%), while 10.9% of heterogeneity was due to phylogeny (Table 4).

The meta-regression models (C and D) used trait category as one of four moderators. The effect size estimates for each trait type were comparable to those from the standard random-effects models (compare Fig. 3A with Fig. 3B or C). Again, the other moderators had no significant effect ($b_{\text{year}} = 0.005$, 95% credible interval, CI $[-0.036, 0.039]$, $b_{\text{Experiment-Observation}} = 0.003$ $[-0.106, 0.116]$, $b_{\text{Field-Laboratory}} = 0.079$ $[-0.074, 0.231]$).

Egger's regression test did not indicate any funnel asymmetry based on meta-analytic residuals from the multilevel meta-regression models (simple: $t_{193} = 1.036$, $P = 0.302$; phylogenetic: $t_{193} = 0.994$, $P = 0.321$; see funnel plots, Fig. 4). However, the trim-and-fill tests indicated that eight (± 8.6 SE; multilevel) and five (± 8.4 SE; phylogenetic) effect sizes were missing, and that we might have overestimated the simple and phylogenetic model means by 0.0093 and 0.0054, respectively. Shifting the CIs for 'avian dataset 1' by these values would result in a credible interval that included zero for the phylogenetic meta-analysis, but not for the simple multilevel meta-analysis. Given this result, and the importance of controlling for phylogeny, our analyses did not provide robust evidence for an offspring sex-ratio bias related to male attractiveness.

675

676 **IV. DISCUSSION**

677 **(1) General findings**

678 We used meta-analysis to test whether females mated to attractive males have a more male-biased
 679 offspring sex ratio (Burley, 1981). There was a small but statistically significant relationship in the
 680 predicted direction. However, male attractiveness accounted for less than 1% of the variation in
 681 offspring sex ratios based on estimates from either standard random-effects meta-analysis or

multilevel meta-analysis. The estimated mean effect sizes were $r = 0.096$ and 0.064 , respectively. Similarly, when we divided the data into different trait categories we found small, but significant, effects of male attractiveness on the sex ratio for two trait types: male ornamentation and male body size. The means for all seven traits were positive, but there were no significant effects for male behavioural displays, body condition, age, extra-pair status or female preference score. Studies measuring ornaments and body size were the most common (these categories contained 37 and 30 studies, respectively, for dataset 1), while there were relatively few studies for any of the other five trait types (6–16 studies; Table 2). The absence of a significant effect for these five trait types might reflect low statistical power. This is plausible given the estimated magnitude of the effects: no matter how the data were analysed, and even when significant, the mean r was always < 0.2 . Alternatively, the lack of significantly positive effect size estimates for some traits might reflect a true null relationship, and/or that unmeasured variation obscured a true relationship. In general though, it is clear that the offspring sex ratio is, on average, only weakly associated with various measures of male attractiveness.

The significant mean effect size observed when combining data from all trait types could be seen as strong evidence for the hypothesis, first proposed by Burley (1981), that there is facultative maternal SRA in response to mate attractiveness/quality. It is, however, prudent to consider the robustness of meta-analytic results to narrow-sense publication bias (i.e. unpublished studies) and/or selective reporting of data (i.e. failure to report non-significant results, or to do so in a way that hinders the ability to calculate effect sizes; see Cassey *et al.*, 2004; review: Jennions *et al.*, 2013). We used increasingly conservative data sets (datasets 2 to 4) to include unknown effect sizes in our analyses, treating them as neutral or opposite to the test hypothesis. Depending on the analysis, the significance of the mean effect tended to vary in its sensitivity to the inclusion of these effects. Specifically, for the standard random effects meta-analysis the estimated mean effect stayed significantly positive for all traits combined across datasets 2, 3 and 4; for ornaments across datasets 2 and 3; and for body size

707 across datasets 2 and 3. For the simple multi-level analysis the estimated mean effect remained
708 significantly positive for all traits combined across avian datasets 1–3.

709 Crucially, the mean effect was not significant for avian dataset 2 when phylogenetic relatedness
710 between species was taken into account (multilevel model B). This analysis is arguably the most
711 important we present. Although it is based only on bird studies, these represent 92% of the available
712 effect sizes. It is appropriate to take phylogeny into account when conducting a meta-analysis that
713 involves reasonably closely related taxa (Chamberlain *et al.*, 2012). There is therefore justifiable
714 concern about the robustness of the main conclusion to selective reporting. In addition, a publication
715 bias test for ‘missing’ studies indicated a strong asymmetry in the funnel plot for all traits. If the
716 putative missing studies were included, the estimated mean effect size was only $r = 0.034$, and did
717 not differ significantly from zero. Again, this suggests that the finding of a significant mean effect
718 size is sensitive to narrow-sense publication bias.

719 Another perspective is that our most important analysis is that based on ornamental male traits. A
720 strict interpretation of the mate attractiveness hypothesis relies on the existence of paternal traits
721 whose heritable benefits are partly male-limited (which is most obviously applicable to ornaments).
722 The initial ‘dataset 1’ analysis indicated a significant effect of ornaments on SRA. At face value, this
723 might reflect the use of traits that are likely to be under the strongest sex-specific selection of any of
724 our categories. A stronger sex bias in the heritable benefits of male traits theoretically leads to
725 stronger SRA results: for example, SRA in response to variation in Fisherian traits (given mutation
726 bias) is predicted to result in more strongly biased offspring sex ratios than is SRA in response to
727 viability indicator traits signalling genes that tend to benefit both sexes (Pen & Weissing, 2000;
728 Fawcett *et al.*, 2007). Again, however, there is cause for concern about publication bias. The
729 significant effect for ornaments was robust to selective reporting (i.e. datasets 2 and 3 still yielded a
730 significant mean effect size), but not to narrow-sense publication bias. A trim-and-fill analysis
731 identified seven ‘missing’ studies that, when imputed, yielded a non-significant mean effect size
732 estimate of $r = 0.029$.

733 The only other trait type with a significant mean effect size was male body size. Why this might be
734 the case is unclear. In birds, which comprise most of our data, there is little evidence that females
735 actively prefer larger males during mate choice. It is noteworthy, however, that body size frequently
736 correlates with major components of male fitness (e.g. success when competing for mates and/or
737 resources, survival, and EPP) in birds (e.g. Kölliker *et al.*, 1999; Kissner, Weatherhead & Francis,
738 2003). More generally, body size often predicts male attractiveness and mating success in other taxa
739 (Fairbairn, Blanckenhorn & Székely, 2007). If sons benefit more than daughters from increased body
740 size then, whether or not females prefer larger males, it could still be adaptive for females to engage
741 in SRA in response to the size of their mate. This is a subtly different explanation from one based on
742 female assessment of mate attractiveness, but only if females do not prefer larger males. It seems
743 possible that the role of male size in attractiveness has been underestimated in birds – or should at
744 least be tested for experimentally. Interestingly, the results for body size are robust to publication
745 bias. The significant effect was not sensitive to selective reporting (i.e. datasets 2 and 3 still yielded a
746 significant mean effect size). It was also robust to narrow-sense publication bias. A trim-and-fill
747 analysis identified seven ‘missing’ studies but their putative values were positive: when imputed, the
748 mean effect size was $r = 0.151$ ($P < 0.001$).

749

750 **(2) Reasons for a small effect: sex allocation decisions are influenced by many factors**

751 There were moderate levels of among-study variation not attributable to sampling error (i.e.
752 heterogeneity) in our data sets (Table 2). This indicates that fairly large differences among studies
753 were unaccounted for. None of our three main moderator variables (year of publication, laboratory
754 *versus* field studies, or observational *versus* experimental studies) had a significant influence on the
755 relationship between male attractiveness and the offspring sex ratio. This was true for both the
756 standard random-effects model and the multilevel analyses. The inclusion of phylogenetic
757 information did little to explain heterogeneity, which remained around 50% (Table 4). The
758 unexplained variance among studies highlights the difficulty of testing specific hypotheses for SRA.

Many selective factors have been suggested to affect SRA and, for some of them, meta-analyses have formally shown that they explain a significant amount of variation in offspring sex ratios (review: West, 2009) (Table 5). These putative forces of selection on SRA are not mutually exclusive; nor, more importantly here, do they act independently of any influence of mate attractiveness on SRA. If multiple sources of selection on SRA interact they could create extremely complex outcomes. This makes it difficult to predict the optimal offspring sex ratio. For example, the benefits of SRA based on mate attractiveness appear to be seasonally dependent (Griffith *et al.*, 2003; Korsten *et al.*, 2006; Delhey *et al.*, 2007), context-dependent (Pryke *et al.*, 2011), or only reliably indicated by specific age classes of males (Taff *et al.*, 2011).

There are numerous other obstacles to testing SRA hypotheses directly. One is the ubiquitous problem of inferring causality from observational studies where there are potential confounding factors that actually generate the observed pattern. For example, assortative mating such that females in good condition tend to mate with more attractive males seems likely in many seasonally breeding bird species. If this occurs, then a male-biased offspring sex ratio might be partly (or entirely) driven by maternal condition (i.e. Trivers & Willard, 1973), rather than by male-biased genetic benefits. Similarly, males might vary in their ability to provide parental care, which could affect SRA (e.g. Bowers *et al.*, 2013). If parental ability is correlated with male attractiveness and sons and daughters differ in the benefits of increased investment, this could lead to adaptive SRA that is correlated with, but not causally related to, attractiveness. Some observational studies in our data set do, however, try to control statistically for confounding maternal and environmental variables (e.g. Addison, Kitaysky & Hipfner, 2008; Taff *et al.*, 2011).

Another potentially common problem arises if the benefits of SRA depend on a male–female interaction. For instance, one of the clearest examples of adaptive SRA – mainly because it involves several different experimental manipulations of male colouration – is in Gouldian finches (*Erythrura gouldiae*). In this species there is severe genetic incompatibility between red and black colour morphs, which has a more detrimental effect on daughters than sons. Females express strong mating

785 preferences for males of the same colour morph. When paired with a male bearing the phenotype of
786 the other morph they exhibit adaptive SRA towards sons (Pryke & Griffith, 2009; Pryke *et al.*, 2011).
787 This species was excluded from our meta-analysis because we only investigated directional female
788 preferences for male traits (i.e. we excluded individual-specific preferences). A type of male–female
789 interaction that might arise in many species and select for SRA based on a non-directional preference
790 is associated with inbreeding. There is evidence that pair relatedness can affect adaptive SRA (e.g.
791 Sardell & DuVal, 2014). If widespread, this could confound patterns of SRA predicted solely on a
792 male’s general attractiveness.

793 Even when genetic incompatibility is absent, SRA might still depend on both female and male
794 qualities. Studies of zebra finches (*Taeniopygia guttata*) suggest that assortative mating accounts for
795 the strong sex ratio biases based on artificial leg colour bands originally detected by Burley (Rutstein
796 *et al.*, 2005). If higher quality females can better bear the costs of producing the more
797 attractive/expensive sex (Katsuki *et al.*, 2012), assortative mating between high-quality females and
798 attractive males could amplify adaptive SRA based solely on male identity. One caveat to this is that
799 the influence of sexually antagonistic genes on adaptive SRA has been poorly explored (but see
800 Calsbeek & Sinervo, 2004; Alonzo & Sinervo, 2007; Blackburn, Albert & Otto, 2010). Given sexual
801 antagonism, it is plausible that assortative mating might eliminate SRA. If genes affecting female
802 quality have more beneficial effects on daughters’ fitness (e.g. Leimar, 1996), while genes that affect
803 male attractiveness more strongly elevate sons’ fitness (Pischedda & Chippindale, 2006), then
804 assortative mating could lead to sex-specific maternal and paternal effects on offspring that nullify
805 the advantages of SRA. If the relative fitness of offspring of each sex remains similar due to
806 ‘symmetrical’ effects of maternal and paternal genes on offspring of the opposite sex (but see
807 Connallon & Jakubowski, 2007; Cox & Calsbeek, 2010), a more equitable sex ratio is predicted. This
808 would occur even if attractive males sire sons of above-average fitness when randomly assigned a
809 mate (i.e. even if there is still a general male-biased benefit of sire attractiveness).

810 Finally, in addition to SRA, there are other, non-mutually exclusive sex allocation responses to the
811 sources of selection described in Table 5, including mate attractiveness. SRA is not the only form of
812 adaptive sex allocation based on changes in investment patterns. There is no evidence in birds that
813 mothers can detect the sex of an embryo and then differentially invest into eggs on this basis (meta-
814 analysis: Rutkowska, Dubiec & Nakagawa, 2014). Mothers can, however, differentially invest into
815 total offspring production (i.e. produce larger broods and/or bigger eggs when mated to an attractive
816 male; Burley, 1986; review: Sheldon, 2000; meta-analysis: Horváthová *et al.*, 2012) and
817 differentially allocate resources towards one sex without any change in the sex ratio or number of
818 offspring per breeding attempt (e.g. Saino *et al.*, 2003). These varied strategic responses are forms of
819 sex allocation that are not always distinguished in the literature, either from each other or from SRA
820 (Carranza, 2002).

821

822 **(3) Reasons for a small effect: measuring sexual selection is tricky**

823 Although there are clear predictions for adaptive SRA based on male attractiveness (West & Sheldon,
824 2002), it is often unclear (*a*) what traits actually determine male attractiveness or quality, and (*b*) if
825 these traits actually affect offspring fitness through male-biased genetic benefits.

826 Identifying targets of mate choice is a perennial problem in comparative analyses or meta-analyses of
827 sexual selection. What makes a male attractive? Many researchers measure several different male
828 traits that are putatively preferred by females. Thus, even within a single empirical study, there can
829 be several reported relationships between ‘attractiveness’ and the offspring sex ratio (e.g. Griffith *et*
830 *al.*, 2003; Bowers *et al.*, 2013). This could be statistically problematic if several male traits are
831 included in a multiple regression as potential predictors of the offspring sex ratio. This was often the
832 case for studies in our data set: many of our effect sizes were taken from generalized linear models
833 (GLMs) or generalized-linear mixed-effects models (GLMMs) with several predictors (exclusion of
834 such data would have halved our sample size). The resultant effect size estimates are therefore the
835 effect of the focal trait controlling for its relationships with other measured traits, and for the

relationship between those traits and the offspring sex ratio (i.e. we reported partial correlations). For a review of the potential statistical biases that occur when using partial rather than bivariate correlations see Aloe (2015), and references therein. The more strongly the measured traits are correlated the greater the discrepancy between partial and bivariate relationships. In general, however, collinearity between measured traits is modest (otherwise they would not be included as independent predictors in a model). A lack of correlation among measured traits does, however, highlight the likelihood that some of them might contribute little if anything to net attractiveness. More generally, female use of multiple cues for mate choice (review: Candolin, 2003) means that effect sizes for individual traits imperfectly capture the true relationship being tested: that between *net male attractiveness* and the offspring sex ratio.

Interpretation of what constitutes an attractive male trait has also expanded since Burley's original hypothesis was formulated. For example, it has been argued that whether a male is a within-pair or extra-pair mate is informative about his relative attractiveness to females (reviews: Westneat & Stewart, 2003; but see Forstmeier *et al.*, 2014). In addition, traits used by females for mate choice are not necessarily those that directly influence adaptive SRA, even if there are male-biased genetic benefits. For example, male attractiveness and competitiveness are often imperfectly correlated (see Kelly, 2008), or even negatively correlated (Hunt *et al.*, 2009). Even if researchers correctly identify heritable traits that determine male attractiveness, these are not necessarily the ones that females assess for the largest potential SRA fitness gains (e.g. if the marginal fitness gains from being more 'dominant' during fights for territories outweigh those of being more attractive to females when holding a territory, and both attractiveness and dominance are heritable).

Determining the relationship between fitness and attractiveness is a major challenge for two reasons. First, it is difficult to measure fitness, and it is often simply assumed that attractiveness predicts net fitness. But do males bearing attractive traits and experiencing higher mating rates actually have greater net fitness than less attractive males? Studies often report an association between male attractiveness/quality and an increase in some components of reproductive success (e.g. number of

862 mates in a single breeding season) without testing whether natural selection (Hine, McGuigan &
863 Blows, 2011) or even other forms of sexual selection (e.g. sperm competitiveness; Lüpold *et al.*,
864 2014) negate the sexually selected benefits of traits that elevate attractiveness. Second, the mate
865 attractiveness hypothesis for adaptive SRA assumes that male-derived fitness benefits exist, and are
866 to some extent sex-limited. In the context of genetic benefits this basically means that male
867 attractiveness is heritable. Many studies have reported significant additive genetic variation for male
868 sexual traits (Prokuda & Roff, 2014), but it remains unclear if net attractiveness (and, ideally, net
869 fitness) is heritable. Genetic covariance among traits affecting fitness can result in the absence of
870 genetic variation in the direction of selection on net attractiveness and, ultimately, fitness (Walsh &
871 Blows, 2009).

872

873 **(4) Updated results and future improvements**

874 Our updated meta-analyses indicate a substantially weaker relationship between measures of mate
875 attractiveness and offspring sex ratios than the $r = 0.205$ reported by West & Sheldon (2002). A
876 decline in effect size over time is a common phenomenon in many research fields, including ecology
877 and evolution (Jennions & Møller, 2002). There are many explanations for a temporal decline,
878 including an early publication bias against weaker effect sizes (i.e. non-significant results) and the
879 inclusion of less appropriate study systems over time (review: Koricheva, Jennions & Lau, 2013).
880 In general, we argue that the observed small effect sizes are expected for many reasons. Six factors
881 that we discussed are: (a) poor control for other selective factors that affect SRA (see Table 5); (b)
882 other forms of conditional sex allocation (e.g. sex-biased provisioning); (c) assortative mating by
883 mate quality; (d) difficulty identifying preferred male traits that directly influence sex allocation –
884 ideally researchers should use measures of *net male attractiveness*; (e) uncertainty that attractiveness
885 increases net male fitness; (f) uncertainty that the benefits of mate choice are genetic and greater for
886 sons than daughters. In addition, there are at least four other factors that might lower effect sizes: (g)
887 high measurement error due to logistic challenges (e.g. measuring behavioural traits accurately); (h)

the inclusion of inappropriate study species (e.g. species with high fecundity are less likely to skew the sex ratio; for a review and explanation see Frank, 1990); (i) difficulty in obtaining appropriate effect size estimates – we had to use data from multiple regression models that included several prospective measures of sire attractiveness because simple bivariate correlations were rarely available; and (j) stochastic noise (e.g. female perceptual errors when assessing males, and constraints on sex-biasing mechanisms).

Aside from considering the problems described above, future studies should broaden their taxonomic scope. Approximately 84% of the studies in our meta-analysis were on birds. This limits the generality of our conclusions, and evaluation of the wider relevance of the SRA mate attractiveness hypothesis. In general, specific sex allocation hypotheses have been tested with a focus on taxa that seem especially amenable (e.g. local mate competition in fig wasps, or helper production in birds and mammals). The disproportionate use of birds to test the SRA mate attractiveness hypothesis might be due to high sexual dimorphism and extravagance of male ornamental traits in many species that suggest strong sexual selection through female choice. More importantly, recent theory suggests that SRA due to mate attractiveness is least likely to occur when there is very strong sexual selection (Fawcett *et al.*, 2011; Booksmythe *et al.*, 2013). Finally, many avian life histories do not conform to key assumptions of mathematical models of sex allocation/adjustment, such as non-overlapping generations, semelparity (or a fixed total investment at every reproductive event), a short and discrete bout of parental investment and investment (potential for control) by one parent only, and no cost of adjustment (review: Komdeur & Pen, 2002). Although some of these limitations will apply to non-avian taxa as well, researchers might wish to reconsider their choice of study species and look to taxa other than birds to test the SRA mate attractiveness hypothesis.

During our attempts to extract effect sizes from papers it became obvious that many studies are guilty of *post hoc* analyses, and fail to clearly specify study goals prior to analysis. Many statistical tests are better seen as exploratory, rather than acting to test a hypothesis with the stated likelihood of type I error (for an approachable case study see Simmons, Nelson & Simonsohn, 2011). There was often an

914 excess of male traits being examined, the inclusion of covariates seemed arbitrary, and the actual
915 statistical test used (e.g. GLM or contingency table) and whether data were pooled or analysed at the
916 brood level was often not justified. All these problems increase the danger of *P*-hacking (selective
917 collection, manipulation, and/or reporting of data to achieve statistical significance) and erroneous
918 reporting of significant results, which inflates effect size estimates (see Simonsohn, Nelson &
919 Simmons, 2014; Head *et al.* 2015). They also made it difficult to locate the information needed to
920 extract effect sizes. The SRA hypothesis is straightforward so it is reasonable to expect greater
921 consistency in how researchers test for a relationship between male attractiveness and offspring sex
922 ratio. We mention this not to criticise colleagues (we are guilty of the same ‘misconduct’ *sensu*
923 Simmons *et al.*, 2011, in some of our own papers), but simply to highlight wider calls by many others
924 (e.g. Garamszegi *et al.*, 2009) for the need to improve standards of reporting in evolutionary ecology
925 studies. It is very possible, for example, that the conclusions of our meta-analysis might have differed
926 if researchers had always used composite measures of male attractiveness (rather than analysing
927 multiple traits) or integrated fitness measures such as female responses, had only controlled for
928 covariates with well-supported biological relevance to their study system and for which they could
929 provide evidence of effects on sex ratios (e.g. level of local mate competition in parasitoids), and had
930 fully reported the direction and magnitude of all tests, regardless of whether or not the effect was
931 significant.

932 Finally, we should note that there is at least one species where well-replicated experimental studies
933 confirm that mothers adjust the offspring sex ratio in response to male phenotype: head colour in
934 Gouldian finches (Pryke & Griffith, 2009; Pryke *et al.*, 2011). As such, there is evidence that SRA in
935 response to male attractiveness has evolved. In our meta-analysis, however, the small mean effect
936 size and its sensitivity to narrow-sense publication bias (the ‘file drawer’ problem, where a greater
937 proportion of non-significant than significant results go unpublished; Rosenthal, 1979; Jennions *et*
938 *al.*, 2013) and/or selective reporting, indicates little support for the hypothesis that such SRA is
939 widespread. Ultimately, the strongest evidence has (and will continue to) come from experiments that

involve direct manipulation of male attractiveness and/or random assignment of males to females. To date there are only a handful of appropriate studies (e.g. Burley, 1981, 1986; Sheldon *et al.*, 1999; Pike & Petrie, 2005; Ferree, 2007). In species where it is feasible, the study design of Sato & Karino (2010), whereby a female's social environment is manipulated so that the same male is assessed to be either relatively attractive or unattractive, seems a particularly powerful approach. Once species in which SRA has been shown robustly to occur are identified, the next step is to test whether this is because sons sired by relatively more attractive males are of higher mean fitness than are daughters. Both lines of evidence are needed to test the SRA mate attractiveness hypothesis fully. Given the many difficulties with the interpretation of observational studies (i.e. inability to predict the sex ratio with certainty due to potentially confounding factors) we suggest that the most fruitful way forward is to conduct experiments that directly rank males according to their mean attractiveness to females, and then randomly assign females a mate of low or high attractiveness. If these experiments are conducted on species where there is little evidence for direct material assistance by males (i.e. no male parental care or mate provisioning) it should be possible to test the hypothesis that genetic benefits that accrue more strongly to sons than daughters can drive greater sex allocation to males.

956

957 **V. CONCLUSIONS**

(1) The mean effect size for the relationship between male attractiveness and offspring sex ratio based on all trait types was small ($r < 0.10$) but significant. Assuming no publication bias this result was consistent using standard random-effects meta-analysis or multilevel meta-analysis models (including a control for phylogenetic relatedness in a data set restricted to birds). (2) These significant mean effect size estimates from all traits combined were, however, sensitive to publication bias – due to a possible ‘file drawer’ effect and/or selective reporting of significant results within studies. Correction for possible bias reduced effect size estimates to non-significant values.

965 (3) The mean effect sizes for seven specific male trait types were significant only for ornamental
966 male traits and measures of male body size. The mean effect remained significant for body size even
967 after correcting for potential publication bias.

968 (4) There was moderate heterogeneity in effect size estimates. The unexplained variation between
969 studies indicates that there are moderating variables that have yet not been identified. We detected no
970 effect of our three test moderators: year of publication, experimental *versus* observational study, and
971 laboratory *versus* field study. There was a very weak phylogenetic signal for birds (i.e. related
972 species did not have similar effect sizes).

973 (5) There was a strong taxonomic bias in our data set. Most studies were on bird species (~ 84%).
974 Given the lack of robust evidence in birds, future studies on other groups (invertebrates, fish, reptiles)
975 are needed to shed light on whether the mate attractiveness SRA hypothesis will have any greater
976 support in other taxa.

977 (6) In sum, our results provide very weak support for the hypothesis that females facultatively adjust
978 the sex ratio of their offspring in response to the attractiveness of their mate.

979

980 **VI. ACKNOWLEDGEMENTS**

981 We thank the following authors who responded to email requests and, where possible, sent data: E.
982 Adkins-Regan, J. Aparicio, R. Bradbury, S. Correa, K. Delhey, A. Dreiss, D. Dowling, P. Dunn, J.
983 Graves, D. Green, L. Gustafsson, M. Hjernquist, A. Husby, P. Korsten, K. Kraaijeveld, D. Leech, S.
984 Leitner, C. Lessells, B. MacDougall-Shackleton, N. Perlut, A. Peters, J. Potti, A. Roulin, T. Schmoll,
985 M. Soma, T. Szekely, C. Taff, D. Westneat, N. Yamaguchi. (Apologies to anyone we have
986 inadvertently omitted.) Funding was provided by the Australian Research Council (M.D.J., J.D.,
987 B.M.), European Research Council grant AdG-294333 to G. Arnqvist (I.B.), the Rutherford
988 Discovery Fellowship, New Zealand (S.N.), and an Australian Research Council Future Fellowship
989 (S.N.).

990

VII. REFERENCES

Studies marked with * contributed effect sizes to the meta-analyses; studies marked with † are key papers from which forward searches were conducted

*ABROE, B., GARVIN, J.C., PEDERSEN, M.C., WHITTINGHAM, L.A. & DUNN, P.O. (2007). Brood sex ratios are related to male size but not to attractiveness in common yellowthroats (*Geothlypis trichas*). *The Auk* **124**, 176-184.

*ADDISON, B., KITAYSKY, A.S. & HIPFNER, J.M. (2008). Sex allocation in a monomorphic seabird with a single-egg clutch: test of the environment, mate quality, and female condition hypotheses. *Behavioral Ecology and Sociobiology* **63**, 135-141.

ALOE, A.M. (2015). Inaccuracy of regression results in replacing bivariate correlations. *Research Synthesis Methods* **6**, 21-27.

ALONZO, S.H. & SINERVO, B. (2007). The effect of sexually antagonistic selection on adaptive sex ratio allocation. *Evolutionary Ecology Research* **9**, 1097-1117.

ANDERSSON, M. (1994). *Sexual Selection*. Princeton University Press, Princeton, NJ.

*APARICIO, J.M. & CORDERO, P.J. (2001). The effects of the minimum threshold condition for breeding on offspring sex-ratio adjustment in the lesser kestrel. *Evolution* **55**, 1188-1197.

BARCLAY, R.M.R. (2012). Variable variation: annual and seasonal changes in offspring sex ratio in a bat. *PLoS ONE* **7**, e36344.

*BELL, S.C., OWENS, I.P.F. & LORD, A.M. (2014). Quality of breeding territory mediates the influence of paternal quality on sex ratio bias in a free-living bird population. *Behavioral Ecology* **25**, 352-358.

*BENITO, M.M., SCHIELZETH, H., GONZÁLEZ-SÓLIS, J. & BECKER, P.H. (2013). Sex ratio adjustments in common terns: influence of mate condition and maternal experience. *Journal of Avian Biology* **44**, 179-188.

1025 BERKELEY, E.V. & LINKLATER, W.L. (2010). Annual and seasonal rainfall may influence progeny sex
1026 ratio in the black rhinoceros. *South African Journal of Wildlife Research* **40**, 53-57.
1027

1028 BLACKBURN, G.S., ALBERT, A.Y.K. & OTTO, S.P. (2010). The evolution of sex ratio adjustment in the
1029 presence of sexually antagonistic selection. *The American Naturalist* **176**, 264-275.
1030

1031 *BLANCHARD, P., HANUISE, N., DANO, S. & WEIMERSKIRCH, H. (2007). Offspring sex ratio in relation
1032 to parental structural size and body condition in the long-lived wandering albatross (*Diomedea*
1033 *exulans*). *Behavioral Ecology and Sociobiology* **61**, 767-773.
1034

1035 BOOKSMYTHE, I., SCHWANZ, L.E. & KOKKO, H. (2013). The complex interplay of sex allocation and
1036 sexual selection. *Evolution* **67**, 673-678.
1037

1038 BORENSTEIN, M., HEDGES, L.V., HIGGINS, J.P.T. & ROTHSTEIN, H.R. (2009). *Introduction to Meta-*
1039 *Analysis*. John Wiley & Sons, Chichester, UK.
1040

1041 *BOWERS, E.K., MUNCLINGER, P., BURES, S., KUCEROVA, L., NADVORNIK, P. & KRIST, M. (2013).
1042 Cross-fostering eggs reveals that female collared flycatchers adjust clutch sex ratios according to
1043 parental ability to invest in offspring. *Molecular Ecology* **22**, 215-228.
1044

1045 BOWERS, E.K., THOMPSON, C.F. & SAKALUK, S.K. (2014). Offspring sex ratio varies with clutch size
1046 for female house wrens induced to lay supernumerary eggs. *Behavioral Ecology* **25**, 165-171.
1047

1048 *BRADBURY, R.B., COTTON, P.A., WRIGHT, J. & GRIFFITHS, R. (1997). Nestling sex ratio in the
1049 European Starling *Sturnus vulgaris*. *Journal of Avian Biology* **28**, 255-258.
1050

1051 BROOKS, R. & KEMP, D.J. (2001). Can older males deliver the good genes? *Trends in Ecology and*
1052 *Evolution* **16**, 308-316.
1053

1054 BROWN, G.R. & SILK, J.B. (2002). Reconsidering the null hypothesis: is maternal rank associated
1055 with birth sex ratios in primate groups? *Proceedings of the National Academy of Sciences of the*
1056 *United States of America* **99**, 11252-11255.
1057

1058 *†BURLEY, N. (1981). Sex-ratio manipulation and selection for attractiveness. *Science* **211**, 721-722.
1059

1060 *BURLEY, N. (1986). Sex-ratio manipulation in color-banded populations of zebra finches. *Evolution*
1061 **40**, 1191-1206.

1062

1063 *CALSBECK, R. & SINERVO, B. (2004). Within-clutch variation in offspring sex determined by
1064 differences in sire body size: cryptic mate choice in the wild. *Journal of Evolutionary Biology* **17**,
1065 464-470.

1066

1067 CAMERON, E.Z. (2004). Facultative adjustment of mammalian sex ratios in support of the Trivers-
1068 Willard hypothesis: evidence for a mechanism. *Proceedings of the Royal Society of London Series B*
1069 **271**, 1723-1728.

1070

1071 CANDOLIN, U. (2003). The use of multiple cues in mate choice. *Biological Reviews* **78**, 575-595.

1072

1073 CARRANZA, J. (2002). What did Trivers and Willard really predict? *Animal Behaviour* **63**, F1-3.

1074

1075 CASSEY, P., EWEN, J.G., BLACKBURN, T.M. & MØLLER, A.P. (2004). A survey of publication bias
1076 within evolutionary ecology. *Proceedings of the Royal Society of London Series B* **271**, S451-454.

1077

1078 CASSEY, P., EWEN, J.G. & MØLLER, A.P. (2006). Revised evidence for facultative sex ratio
1079 adjustment in birds: a correction. *Proceedings of the Royal Society of London Series B* **273**, 3129-
1080 3130.

1081

1082 CHAMBERLAIN, S.A., HOVICK, S.M., DIBBLE, C.J., RASMUSSEN, N.L., VAN ALLEN, B.G., MAITNER,
1083 B.S., AHERN, J.R., BELL-DERESKE, L.P., ROY, C.L., MEZA-LOPEZ, M., CARRILLO, J., SIEMANN, E.,
1084 LAJEUNESSE, M.J. & WHITNEY, K.D. (2012). Does phylogeny matter? Assessing the impact of
1085 phylogenetic information in ecological meta-analysis. *Ecology Letters* **15**, 627-636.

1086

1087 CHARNOV, E.L. (1979). The genetical evolution of patterns of sexuality: Darwinian fitness. *The*
1088 *American Naturalist* **113**, 465-480.

1089

1090 CICHON, M., DUBIEC, A. & STOCZKO, M. (2003). Laying order and offspring sex in blue tits *Parus*
1091 *caeruleus*. *Journal of Avian Biology* **34**, 355-359.

1092

1093 CLUTTON-BROCK, T.H. (1986). Sex ratio variation in birds. *Ibis* **128**, 317-329.

1094

1095 CLUTTON-BROCK, T.H., ALBON, S.D. & GUINNESS, F.E. (1984). Maternal dominance, breeding
1096 success and birth sex-ratios in red deer. *Nature* **308**, 358-360.

1097

1098 *COCKBURN, A. & DOUBLE, M.C. (2008). Cooperatively breeding superb fairy-wrens show no
1099 facultative manipulation of offspring sex ratio despite plausible benefits. *Behavioral Ecology and*
1100 *Sociobiology* **62**, 681-688.

1101

1102 COCKBURN, A., SCOTT, M.P. & DICKMAN, C.R. (1985). Sex-ratio and intrasexual kin competition in
1103 mammals. *Oecologia* **66**, 427-429.

1104

1105 COHEN, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*, 2nd Edition. Lawrence
1106 Erlbaum, Hillsdale, NJ.

1107

1108 CONNALLON, T. & CLARK A.G. (2014) Evolutionary inevitability of sexual antagonism. *Proceedings*
1109 *of the Royal Society of London Series B* **281**, 20132013.

1110

1111 CONNALLON, T. & JAKUBOWSKI, E. (2007). Association between sex ratio distortion and sexually
1112 antagonistic fitness consequences of female choice. *Evolution* **63**, 2179-2183.

1113

1114 *CORREA, S.M., HORAN, C.M., JOHNSON, P.A. & ADKINS-REGAN, E. (2011). Copulatory behaviors
1115 and body condition predict post-mating female hormone concentrations, fertilization success, and
1116 primary sex ratios in Japanese quail. *Hormones and Behavior* **59**, 556-564.

1117

1118 *COTTON, A.J., FÖLDIVARI, M., COTTON, S. & POMIANKOWSKI, A. (2014). Male eyespan size is
1119 associated with meiotic drive in wild stalk-eyed flies (*Teleopsis dalmanni*). *Heredity* **112**, 363-369.

1120

1121 COTTON, S., FOWLER, K. & POMIANKOWSKI, A. (2004). Do sexual ornaments exhibit heightened
1122 condition-dependent expression as predicted by the handicap hypothesis? *Proceedings of the Royal*
1123 *Society of London Series B* **271**, 771-783.

1124

1125 COX, R.M. & CALSBEEK, R. (2010). Cryptic sex-ratio bias provides indirect genetic benefits despite
1126 sexual conflict. *Science* **328**, 92-94.

1127

1128 *COX, R.M., DURYEA, M.C., NAJARRO, M. & CALSBEEK, R. (2011). Paternal condition drives
1129 progeny sex-ratio bias in a lizard that lacks parental care. *Evolution* **65**, 220-230.

1130

1131 CRUICKSHANK, T. & WADE, M.J. (2012). Maternal adjustment of the sex ratio in broods of the broad-
1132 horned flour beetle, *Gnathocerus cornutus*. *Integrative and Comparative Biology* **52**, 100-107.

1133

1134 *CUNNINGHAM, E. J. A. AND A. F. RUSSELL (2001). Sex differences in avian yolk hormone levels -
1135 Reply. *Nature* **412**, 498-499.

1136

1137 *DEBIAK, A.L., MCCORMICK, D.L., KAPLAN, J.D., TISCHLER, K.B. & LINDSAY, A.R. (2014). A
1138 molecular genetic assessment of sex ratios from pre-fledged juvenile and migrating adult common
1139 loons (*Gavia immer*). *Waterbirds* **37**, 6-15.

1140

1141 *DELHEY, K., PETERS, A., JOHNSEN, A. & KEMPENAERS, B. (2007). Brood sex ratio and male UV
1142 ornamentation in blue tits (*Cyanistes caeruleus*): correlational evidence and an experimental test.
1143 *Behavioral Ecology and Sociobiology* **61**, 853-862.

1144

1145 *DELMORE, K.E., KLEVEN, O., LASKEMOEN, T., CROWE, S.A., LIFJELD, J.T. & ROBERTSON, R.J.
1146 (2008). Sex allocation and parental quality in tree swallows. *Behavioral Ecology* **19**, 1243-1249.

1147

1148 *DIETRICH-BISCHOFF, V., SCHMOLL, T., WINKEL, W., KRACKOW, S. & LUBJUHNS, T. (2006). Extra-pair
1149 paternity, offspring mortality and offspring sex ratio in the socially monogamous coal tit (*Parus*
1150 *ater*). *Behavioral Ecology and Sociobiology* **60**, 563-571.

1151

1152 *DOLAN, A.C., MURPHY, M.T., REDMOND, L.J. & DUFFIELD, D. (2009). Maternal characteristics and
1153 the production and recruitment of sons in the eastern kingbird (*Tyrannus tyrannus*). *Behavioral*
1154 *Ecology and Sociobiology* **63**, 1527-1537.

1155

1156 *DOWLING, D.K. & MULDER, R.A. (2006). Combined influence of maternal and paternal quality on
1157 sex allocation in red-capped robins. *Journal of Evolutionary Biology* **19**, 440-449.

1158

1159 *DREISS, A., RICHARD, M., MOYEN, F., WHITE, J., MØLLER, A.P. & DANCHIN, E. (2006). Sex ratio and
1160 male sexual characters in a population of blue tits, *Parus caeruleus*. *Behavioral Ecology* **17**, 13-19.

1161

1162 *DRULLION, D. & DUBOIS, F. (2011). Neighbours' breeding success and the sex ratio of their
1163 offspring affect the mate preferences of female zebra finches. *PLoS One* **6**, e29737.

1164

1165 *DU, B. & LU, X. (2010). Sex allocation and paternity in a cooperatively breeding passerine:
 1166 evidence for the male attractiveness hypothesis? *Behavioral Ecology and Sociobiology* **64**, 1631-
 1167 1639.

1168

1169 DUBOIS, N.S., KENNEDY, E.D. & GETTY, T. (2006). Surplus nest boxes and the potential for polygyny
 1170 affect clutch size and offspring sex ratio in house wrens. *Proceedings of the Royal Society of London*
 1171 *Series B* **273**, 1751-1757.

1172

1173 DUNN, P.O., WHITTINGHAM, L.A. & PITCHER, T.E. (2001). Mating systems, sperm competition, and
 1174 the evolution of sexual dimorphism in birds. *Evolution* **55**, 161-175.

1175

1176 DÜSING, C. (1884). *Die Regulierung des Geschlechtsverhältnisses bei der Vermehrung der*
 1177 *Menschen, Tiere und Pflanzen*. Fischer, Jena, Germany.

1178

1179 DUVAL, S. & TWEEDIE, R. (2000a). A nonparametric "trim and fill" method of accounting for
 1180 publication bias in meta-analysis. *Journal of the American Statistical Association* **95**, 89-98.

1181

1182 DUVAL, S. & TWEEDIE, R. (2000b). Trim and fill: A simple funnel-plot-based method of testing and
 1183 adjusting for publication bias in meta-analysis. *Biometrics* **56**, 455-463.

1184

1185 EDVARDSSON, M. & ARNQVIST, G. (2005). The effects of copulatory courtship on differential
 1186 allocation in the red flour beetle *Tribolium castaneum*. *Journal of Insect Behavior* **18**, 313-322.

1187

1188 EGGER, M., DAVEY SMITH, G., SCHNEIDER, M. & MINDER, C. (1997). Bias in meta-analysis detected
 1189 by a simple, graphical test. *British Medical Journal* **315**, 629-634.

1190

1191 *†ELLEGGREN, H., GUSTAFSSON, L. & SHELDON, B.C. (1996). Sex ratio adjustment in relation to
 1192 paternal attractiveness in a wild bird population. *Proceedings of the National Academy of Sciences of*
 1193 *the United States of America* **93**, 11723-11728.

1194

1195 EWEN, J.G., CASSEY, P. & MØLLER, A.P. (2004). Facultative primary sex ratio variation: a lack of
 1196 evidence in birds? *Proceedings of the Royal Society of London Series B* **271**, 1277-1282.

1197

1198 FAIRBAIRN, D.J., BLANCKENHORN, W.U. & SZÉKELY, T. (eds) (2007). *Sex, Size and Gender Roles:*
 1199 *evolutionary studies of sexual size dimorphism*. Oxford University Press, Oxford, UK.

1200

1201 *FARGALLO, J.A., DÁVILA, J.A., POTTI, J., DE LEÓN, A. & POLO, V. (2004). Nest size and hatchling
 1202 sex ratio in chinstrap penguins. *Polar Biology* **27**, 339-343.

1203

1204 FARRIS, H.E. & RYAN, M.J. (2011). Relative comparisons of call parameters enable auditory grouping
 1205 in frogs. *Nature Communications* **2**, 410.

1206

1207 †FAWCETT, T.W., KUIJPER, B., PEN, I. & WEISSING, F.J. (2007). Should attractive males have more
 1208 sons? *Behavioral Ecology* **18**, 71-80.

1209

1210 FAWCETT, T.W., KUIJPER, B., WEISSING, F.J. & PEN, I. (2011). Sex-ratio control erodes sexual
 1211 selection, revealing evolutionary feedback from adaptive plasticity. *Proceedings of the National*
 1212 *Academy of Sciences of the United States of America* **108**, 15925-15930.

1213

1214 FIRMAN, R.C. (2011). Polyandrous females benefit by producing sons that achieve high reproductive
 1215 success in a competitive environment. *Proceedings of the Royal Society of London Series B* **278**,
 1216 2823-2831.

1217

1218 FISHER, R.A. (1930). *The Genetical Theory of Natural Selection*. Clarendon, Oxford, UK.

1219

1220 *FERREE, E.D. (2007). White tail plumage and brood sex ratio in dark-eyed juncos (*Junco hyemalis*
 1221 *thurberi*). *Behavioral Ecology and Sociobiology* **62**, 109-117.

1222

1223 FORSTMEIER, W., NAKAGAWA, S., GRIFFITH, S.C. & KEMPENAERS, B. (2014). Female extra-pair
 1224 mating: adaptation or genetic constraint? *Trends in Ecology and Evolution* **29**, 456-464.

1225

1226 †FRANK, S.A. (1990). Sex allocation theory for birds and mammals. *Annual Review of Ecology and*
 1227 *Systematics* **21**, 13-55.

1228

1229 FRANK, S.A. (1998). *Foundations of Social Evolution*. Princeton University Press, Princeton, N.J.

1230

1231 GARAMSZEGI, L.Z., CALHIM, S., DOCHTERMANN, N.A., HEGYI, G., HURD, P.L., JØREGENSEN, C.,
 1232 KUTSUKAKE, N., LAJEUNESSE, M.J., POLLATD, K.A., SCHIELZETH, H., SYMONDS, M.R.E. &
 1233 NAKAGAWA, S. (2009). Changing philosophies and tools for statistical inferences in behavioral
 1234 ecology. *Behavioral Ecology* **20**, 1363-1375.

1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269

- GILDERSLEEVE, K., HASELTON, M.G. & FALES, M.R. (2014). Do women's mate preferences change across the ovulatory cycle? A meta-analytic review. *Psychological Bulletin* **140**, 1205-1259.
- GODFRAY, H.C.J. (1994). *Parasitoids: behavioural and evolutionary ecology* pp. 151-210. Princeton University Press, Princeton, NJ.
- GOWATY, P.A. & LENNARTZ, M.R. (1985). Sex-ratios of nestling and fledgling red-cockaded woodpeckers (*Picoides borealis*) favor males. *The American Naturalist* **126**, 347-353.
- GRAHAM, E.B., CARO, S.P. & SOCKMAN, K.W. (2011). Change in offspring sex ratio over a very short season in Lincoln's sparrows: the potential role of bill development. *Journal of Field Ornithology* **82**, 44-51.
- GRANA, S.C., SAKALUK, S.K., BOWDEN, R.M., DOELLMAN, M.A., VOGEL, L.A. & THOMPSON, C.F. (2012). Reproductive allocation in female house wrens is not influenced by experimentally altered male attractiveness. *Behavioral Ecology and Sociobiology* **66**, 1247-1258.
- *GREEN, D.J. (2002). Pair bond duration influences paternal provisioning and the primary sex ratio of brown thornbill broods. *Animal Behaviour* **64**, 791-800.
- GRIFFIN, A.S., SHELDON, B.C. & WEST, S.A. (2005). Cooperative breeders adjust offspring sex ratios to produce helpful helpers. *The American Naturalist* **166**, 628-632.
- *GRIFFITH, S.C., ÖRNBORG, J., RUSSELL, A.F., ANDERSSON, S. & SHELDON, B.C. (2003). Correlations between ultraviolet coloration, overwinter survival and offspring sex ratio in the blue tit. *Journal of Evolutionary Biology* **16**, 1045-1054.
- *GRINDSTAFF, J.L., BUERKLE, C.A., CASTO, J.M., NOLAN JR., V. & KETTERSON, E.D. (2001). Offspring sex ratio is unrelated to male attractiveness in dark-eyed juncos (*Junco hyemalis*). *Behavioral Ecology and Sociobiology* **50**, 312-316.
- HADFIELD, J.D. (2010). MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software* **33**, 1-22.

1270 HADFIELD, J.D. & NAKAGAWA, S. (2010). General quantitative genetic methods for comparative
1271 biology: phylogenies, taxonomies and multi-trait models for continuous and categorical characters.
1272 *Journal of Evolutionary Biology* **23**, 494-508.
1273

1274 HAMILTON, W.D. (1967). Extraordinary sex ratios. *Science* **156**, 477-488.
1275

1276 HEAD, M.L., HOLMAN, L., LANFEAR, R., KAHN, A.T. & JENNIONS, M.D. (2015). The extent and
1277 consequences of P-hacking in science. *Public Library of Science, Biology* **13**: e1002106.
1278

1279 HEDGES, L. & OLKIN, I. (1985). *Statistical methods for meta-analysis*. Academic Press, New York,
1280 NY.
1281

1282 *HELLE, S., LAAKSONEN, T., ADAMSSON, A., PARANKO, J. & HUITU, O. (2008). Female field voles
1283 with high testosterone and glucose levels produce male-biased litters. *Animal Behaviour* **75**, 1031-
1284 1039.
1285

1286 HIGGINS, J.P.T., THOMPSON, S.G., DEEKS, J.J. & ALTMAN, D.G. (2003). Measuring inconsistency in
1287 meta-analyses. *British Medical Journal* **327**, 557-560.
1288

1289 HINE, E., MCGUIGAN, K. & BLOWS, M.W. (2011). Natural selection stops the evolution of male
1290 attractiveness. *Proceedings of the National Academy of Sciences of the United States of America* **108**,
1291 3659-3664
1292

1293 *HJERNQUIST, M.B., HJERNQUIST, K.A.T., FORSMAN, J.T. & GUSTAFSSON, L. (2009). Sex allocation
1294 in response to local resource competition over breeding territories. *Behavioral Ecology* **20**, 335-339.
1295

1296 HOELZER, G.A. (1989). The good parent process of sexual selection. *Animal Behaviour* **38**, 1067-
1297 1078.
1298

1299 HOLMAN, L. & KOKKO, H. (2013). The consequences of polyandry for population viability, extinction
1300 risk and conservation. *Philosophical Transactions of the Royal Society of London Series B* **368**,
1301 20120053.
1302

1303 HORVÁTHOVÁ, T., NAKAGAWA, S. & ULLER, T. (2012). Strategic female reproductive investment in
1304 response to male attractiveness in birds. *Proceedings of the Royal Society of London Series B* **279**,
1305 163-170.

1306

1307 *HOUSE, C.M., SIMMONS, L.W., KOTIAHO, J.S., TOMKINS, J.L. & HUNT, J. (2011). Sex ratio bias in
1308 the dung beetle *Onthophagus taurus*: adaptive allocation or sex-specific offspring mortality?
1309 *Evolutionary Ecology* **25**, 363-372.

1310

1311 HOUSWORTH, E.A., MARTINS, E.P. & LYNCH, M. (2004). The phylogenetic mixed model. *The*
1312 *American Naturalist* **163**, 84-96.

1313

1314 HUNT, J., BREUKER, C.J., SADOWSKI, J.A. & MOORE, A.J. (2009). Male-male competition, female
1315 mate choice and their interaction: determining total sexual selection. *Journal of Evolutionary*
1316 *Biology* **22**, 13-26.

1317

1318 *HUSBY, A., SAETHER, B.E., JENSEN, H. & RINGSBY, T.H. (2006). Causes and consequences of
1319 adaptive seasonal sex ratio variation in house sparrows. *Journal of Animal Ecology* **75**, 1128-1139.

1320

1321 JANSSEN, K., ERIKSTAD, K.E. & BENSCH, S. (2006). Offspring sex ratio allocation in the parasitic
1322 jaeger: selection for pale females and melanic males? *Behavioral Ecology* **17**, 236-245.

1323

1324 JENNIONS, M.D. & MØLLER, A.P. (2002). Relationships fade with time: a meta-analysis of temporal
1325 trends in publication in ecology and evolution. *Proceedings of the Royal Society of London Series B*
1326 **269**, 43-48.

1327

1328 JENNIONS, M.D. & MØLLER, A.P. (2003). A survey of the statistical power of research in behavioral
1329 ecology and animal behavior. *Behavioral Ecology* **14**, 438-445.

1330

1331 JENNIONS, M.D., KAHN, A.T., KELLY, C.D. & KOKKO, H. (2012). Meta-analysis and sexual selection:
1332 past studies and future possibilities. *Evolutionary Ecology* **26**, 1119-1151.

1333

1334 JENNIONS, M.D., LORTIE, C.J., ROSENBERG, M.S. & ROTHSTEIN, H.R. (2013). Publication and related
1335 biases. In: *Handbook of Meta-Analysis in Ecology and Evolution* (eds. J. KORICHEVA, J. GUREVITCH
1336 & K. MENGENSEN), pp. 207-236. Princeton University Press, Princeton, NJ.

1337

1338 JETZ, W., THOMAS, G.H., JOY, J.B., HARTMANN, K. & MOOERS, A.O. (2012). The global diversity of
1339 birds in space and time. *Nature* **491**, 444-448.

1340

1341 *JOHNSON, L.S., THOMPSON, C.F., SAKALUK, S.K., NEUHÄUSER, M., JOHNSON, B.G.P., SOUKUP, S.S.,
1342 FORSYTHE, S.J. & MASTERS, B.S. (2009). Extra-pair young in house wren broods are more likely to
1343 be male than female. *Proceedings of the Royal Society of London Series B* **276**, 2285-2289.

1344

1345 JONES, K.S., NAKAGAWA, S. & SHELDON, B.C. (2009). Environmental sensitivity in relation to size
1346 and sex in birds: meta-regression analysis. *The American Naturalist* **174**, 122-133.

1347

1348 *KARINO, K., KOBAYASHI, M. & ORITA, K. (2006). Adaptive offspring sex ratio depends on male tail
1349 length in the guppy. *Ethology* **112**, 1050-1055.

1350

1351 *KARINO, K. & SATO, A. (2009). Male-biased sex ratios in offspring of attractive males in the guppy.
1352 *Ethology* **115**, 682-690.

1353

1354 *KATSUKI, M., HARANO, T., MIYATAKE, T., OKADA, K. & HOSKEN, D. (2012). Intralocus sexual
1355 conflict and offspring sex ratio. *Ecology Letters* **15**, 193-197.

1356

1357 KELLY, C.D. (2008). The interrelationships between resource-holding potential, resource-value and
1358 reproductive success in territorial males: how much variation can we explain? *Behavioral Ecology*
1359 *and Sociobiology* **62**, 855-871.

1360

1361 *KEMPENAERS, B., VERHEYEN, G.R. & DHONDT, A.A. (1997). Extrapair paternity in the blue tit
1362 (*Parus caeruleus*): female choice, male characteristics, and offspring quality. *Behavioral Ecology* **8**,
1363 481-492.

1364

1365 KIE, J.G., JOHNSON, B.K., NOYES, J.H., WILLIAMS, C.L., DICK, B.L., RHODES, O.E., STUSSY, R.J. &
1366 BOWYER, R.T. (2013). Reproduction in North American elk *Cervus elaphus*: paternity of calves sired
1367 by males of mixed age classes. *Wildlife Biology* **19**, 302-310.

1368

1369 KISSNER, K.J., WEATHERHEAD, P.J. & FRANCIS, C.M. (2003). Sexual size dimorphism and timing of
1370 spring migration in birds. *Journal of Evolutionary Biology* **16**, 154-162.

1371

1372 *KÖLLIKER, M., HEEB, P., WERNER, I., MATEMAN, A.C., LESSELLS, C.M. & RICHNER, H. (1999).
1373 Offspring sex ratio is related to male body size in the great tit (*Parus major*). *Behavioral Ecology* **10**,
1374 68-72.

1375

1376 KOMDEUR, J. & PEN, I. (2002). Adaptive sex allocation in birds: the complexities of linking theory
1377 and practice. *Philosophical Transactions of the Royal Society of London Series B* **357**, 373-380.

1378

1379 KORICHEVA, J., JENNIONS, M.D. & LAU, J. (2013). Temporal trends in effect sizes: causes, detection
1380 and implication. In: *Handbook of Meta-Analysis in Ecology and Evolution* (eds J. Koricheva, J.
1381 Gurevitch & K. Mengersen), pp. 237-254. Princeton University Press, Princeton, NJ.

1382

1383 *KORPIMÄKI, E., MAY, C.A., PARKIN, D.T., WETTON, J.H. & WIEHN, J. (2000). Environmental- and
1384 parental condition-related variation in sex ratio of kestrel broods. *Journal of Avian Biology* **31**, 128-
1385 134.

1386

1387 *KORSTEN, P., LESSELLS, C.M., MATEMAN, A.C., VAN DER VELDE, M. & KOMDEUR, J. (2006).
1388 Primary sex ratio adjustment to experimentally reduced male UV attractiveness in blue tits.
1389 *Behavioral Ecology* **17**, 539-546.

1390

1391 *KRAAIJEVELD, K., MING, M., KOMDEUR, J. & MULDER, R.A. (2007). Offspring sex ratios in relation
1392 to mutual ornamentation and extra-pair paternity in the Black Swan *Cygnus atratus*. *Ibis* **149**, 79-85.

1393

1394 KRACKOW, S. (1995). Potential mechanisms for sex ratio adjustment in mammals and birds.
1395 *Biological Reviews* **70**, 225-241.

1396

1397 KRUUK, L.E.B., CLUTTON-BROCK, T.H., ALBON, S.D., PEMBERTON, J.M. & GUINNESS, F.E. (1999).
1398 Population density affects sex ratio variation in red deer. *Nature* **399**, 459-461.

1399

1400 *LEE, J.-W. & HATCHWELL, B.J. (2011). Do females adjust brood sex ratio according to males'
1401 genetic structures in a gregarious passerine, the vinous-throated parrotbill *Paradoxornis webbianus*?
1402 *Journal of Avian Biology* **42**, 447-453.

1403

1404 *LEECH, D.I., HARTLEY, I.R., STEWART, I.R.K., GRIFFITH, S.C. & BURKE, T. (2001). No effect of
1405 parental quality or extrapair paternity on brood sex ratio in the blue tit (*Parus caeruleus*). *Behavioral*
1406 *Ecology* **12**, 674-680.

1407

1408 LEIMAR, O. (1996). Life-history analysis of the Trivers and Willard sex-ratio problem. *Behavioral*
1409 *Ecology* **7**, 316-325.

1410

1411 *LEITNER, S., MARSHALL, R.C., LEISLER, B. & CATCHPOLE, C.K. (2006). Male song quality, egg size
1412 and offspring sex in captive canaries (*Serinus canaria*). *Ethology* **112**, 554-563.

1413

1414 *LEONARD, M.L. & WEATHERHEAD, P.J. (1996). Dominance rank and offspring sex ratios in
1415 domestic fowl. *Animal Behaviour* **51**, 725-731.

1416

1417 *LESSELLS, C.M., MATEMAN, A.C. & VISSER, J. (1996). Great tit hatchling sex ratios. *Journal of*
1418 *Avian Biology* **27**, 135-142.

1419

1420 LEZALOVA, R., TKADLEK, E., OBORNIK, M., SIMEK, J. & HONZA, M. (2005). Should males come first?
1421 The relationship between offspring hatching order and sex in the black-headed gull *Larus ridibundus*.
1422 *Journal of Avian Biology* **36**, 478-483.

1423

1424 LIMBOURG, T., MATEMAN, A.C., ANDERSSON, S. & LESSELLS, C.M. (2004). Female blue tits adjust
1425 parental effort to manipulated male UV attractiveness. *Proceedings of the Royal Society of London*
1426 *Series B* **271**, 1903-1908.

1427

1428 *LIMBOURG, T., MATEMAN, A.C. & LESSELLS, C.M. (2013). Parental care and UV coloration in blue
1429 tits: opposite correlations in males and females between provisioning rate and mate's coloration.
1430 *Journal of Avian Biology* **44**, 17-26.

1431

1432 LIPSEY, M.W. & WILSON, D.B. (2001). *Practical meta-analysis*. Sage Publishing, Beverly Hills, CA.

1433

1434 LISLEVAND, T., BYRKJEDAL, I., BORGE, T. & SAETRE, G.P. (2005). Egg size in relation to sex of
1435 embryo, brood sex ratios and laying sequence in northern lapwings (*Vanellus vanellus*). *Journal of*
1436 *Zoology* **267**, 81-87.

1437

1438 *LONG, T.A.F. & PISCHEDDA, A. (2005). Do female *Drosophila melanogaster* adaptively bias
1439 offspring sex ratios in relation to the age of their mate? *Proceedings of the Royal Society of London*
1440 *Series B* **272**, 1781-1787.

1441

1442 LOYAU, A. & LACROIX, F. (2010). Watching sexy displays improves hatching success and offspring
1443 growth through maternal allocation. *Proceedings of the Royal Society of London Series B* **277**, 3453-
1444 3460.

1445

1446 LOYAU, A., SAINT JALME, M., MAUGET, R. & SORCI, G. (2007). Male sexual attractiveness affects the
1447 investment of maternal resources into the eggs in peafowl (*Pavo cristatus*). *Behavioral Ecology and*
1448 *Sociobiology* **61**, 1043-1052.

1449

1450 *LU, X., ZENG, X. & DU, B. (2013). Body attributes of both parents jointly affect offspring sex
1451 allocation in a socially monogamous, size-monomorphic passerine. *Current Zoology* **59**, 271-277.

1452

1453 LÜPOLD, S., TOMKINS, J.L., SIMMONS, L.W. & FITZPATRICK, J.L. (2014). Female monopolization
1454 mediates the relationship between pre- and post-copulatory sexual traits. *Nature Communications* **5**,
1455 3184.

1456

1457 MAESTRIPIERI, D. (2002). Maternal dominance rank and age affect offspring sex ratio in pigtail
1458 macaques. *Journal of Mammalogy* **83**, 563-568.

1459

1460 *MAGRATH, M.J.L., GREEN, D.J. & KOMDEUR, J. (2002). Sex allocation in the sexually monomorphic
1461 fairy martin. *Journal of Avian Biology* **33**, 260-268.

1462

1463 MENGENSEN, K., JENNIONS, M.D. & SCHMID, C.H. (2013). Statistical models for the meta-analysis of
1464 nonindependent data. In: *Handbook of Meta-Analysis for Ecology and Evolution* (eds. J. KORICHEVA,
1465 J. GUREVITCH & K. MENGENSEN), pp. 255-283. Princeton University Press, Princeton, NJ.

1466

1467 MEUNIER, J., WEST, S.A. & CHAPUISAT, M. (2008). Split sex ratios in the social Hymenoptera: a
1468 meta-analysis. *Behavioral Ecology* **19**, 382-390.

1469

1470 MØLLER, A.P. & BIRKHEAD, T. (1994). The evolution of plumage brightness in birds is related to
1471 extrapair paternity. *Evolution* **48**, 1089-1100.

1472

1473 MØLLER, A.P. & JENNIONS, M.D. (2001). How important are direct fitness benefits of sexual
1474 selection? *Naturwissenschaften* **88**, 401-415.

1475

1476 MØLLER, A.P. & NINNI, P. (1998). Sperm competition and sexual selection: a meta-analysis of
1477 paternity studies of birds. *Behavioral Ecology and Sociobiology* **43**, 345-358.
1478

1479 MOORE, A.J., GOWATY, P.A., WALLIN, W.G. & MOORE, P.J. (2001). Sexual conflict and the evolution
1480 of female mate choice and male social dominance. *Proceedings of the Royal Society of London Series*
1481 *B* **268**, 517-523.
1482

1483 MURTAUGH, P.A. (2002). Journal quality, effect size, and publication bias in meta-analysis. *Ecology*
1484 **83**, 1162-1166.
1485

1486 NAKAGAWA, S. & CUTHILL, I.C. (2007). Effect size, confidence interval and statistical significance: a
1487 practical guide for biologists. *Biological Reviews* **82**, 591-605.
1488

1489 NAKAGAWA, S. & SANTOS, E.S.A. (2012). Methodological issues and advances in biological meta-
1490 analysis. *Evolutionary Ecology*, **26**, 1253-1274.
1491

1492 *NETO, J.M., HANSSON, B. & HASSELQUIST, D. (2011). Sex allocation in Savi's warblers *Locustella*
1493 *luscinioides*: multiple factors affect seasonal trends in brood sex ratios. *Behavioral Ecology and*
1494 *Sociobiology* **65**, 297-304.
1495

1496 NEVILLE, K.J., VALLENDER, R. & ROBERTSON, R.J. (2008). Nestling sex ratio of golden-winged
1497 warblers *Vermivora chrysoptera* in an introgressed population. *Journal of Avian Biology* **39**, 599-
1498 604.
1499

1500 NEVISON, C. (1997). Social rank and birth sex ratios in rhesus monkeys (*Macaca mulatta*). *Applied*
1501 *Animal Behaviour Science* **51**, 285-291.
1502

1503 *NISHIUMI, I. (1998). Brood sex ratio is dependent on female mating status in polygynous great reed
1504 warblers. *Behavioral Ecology and Sociobiology* **44**, 9-14.
1505

1506 *ODDIE, K.R. & REIM, C. (2002). Egg sex ratio and paternal traits: using within-individual
1507 comparisons. *Behavioral Ecology* **13**, 503-510.
1508

1509 OLSSON, M., MADSEN, T., ULLER, T., WAPSTRA, E. & UJVARI, B. (2005a). The role of Haldane's rule
1510 in sex allocation. *Evolution* **59**, 221-225.

1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545

*OLSSON, M., WAPSTRA, E. & ULLER, T. (2005*b*). Differential sex allocation in sand lizards: bright males induce daughter production in a species with heteromorphic sex chromosomes. *Biology Letters* **1**, 378-380.

PALMER, A.R. (2000). Quasireplication and the contract of error: lessons from sex ratios, heritabilities and fluctuating asymmetry. *Annual Review of Ecology and Systematics* **31**, 441-480.

*PARISER, E.C., GILBERT, L., HAZON, N., ARNOLD, K.E. & GRAVES, J.E. (2012). Mind the gap: the ratio of yolk androgens and antioxidants varies between sons and daughters dependent on paternal attractiveness. *Behavioral Ecology and Sociobiology* **66**, 519-527.

*PARKER, T.H. (2005). No evidence for adaptive differential sex allocation in red junglefowl (*Gallus gallus*). *The Auk* **122**, 1161-1168.

PARKER, T.H. (2013). What do we really know about the signaling role of plumage colour in blue tits? A case study of impediments to progress in evolutionary biology. *Biological Reviews* **88**, 511-536.

PEN, I. & WEISSING, F.J. (2000). Sexual selection and the sex ratio: an ESS analysis. *Selection* **1**, 111-121.

PEREIRA, T.V. & IOANNIDIS, J.P.A. (2011). Statistically significant meta-analyses of clinical trials have modest credibility and inflated effects. *Journal of Clinical Epidemiology* **64**, 1060-1069.

*PERLUT, N.G., TRAVIS, S.E., DUNBAR, C.A., STRONG, A.M. & WRIGHT, D.M. (2014). Nestling sex ratios do not support long-term parity in two species with different life-history strategies. *The Auk* **131**, 224-234.

PETRIE, M., SCHWABL, H., BRANDE-LAVRIDSEN, N. & BURKE, T. (2001). Sex differences in avian yolk hormone levels. *Nature* **412**, 498.

*PIKE, T.W. & PETRIE, M. (2005). Offspring sex ratio is related to paternal train elaboration and yolk corticosterone in peafowl. *Biology Letters* **1**, 204-207.

1546 PISCHEDDA, A. & CHIPPINDALE, A. (2006). Intra-locus sexual conflict diminishes the benefits of
1547 sexual selection. *Public Library of Science Biology* **4**, 2099-2103.

1548

1549 *POLO, V., VEIGA, J.P., CORDERO, P.J., VIÑUELA, J. & MONAGHAN, P. (2004). Female starlings adjust
1550 primary sex ratio in response to aromatic plants in the nest. *Proceedings of the Royal Society of*
1551 *London Series B* **271**, 1929-1933.

1552

1553 POMIANKOWSKI, A., IWASA, Y. & NEE, S. (1991). The evolution of costly mate preferences I. Fisher
1554 and biased mutation. *Evolution* **45**, 1422-1430.

1555

1556 *POTVIN, D.A. & MACDOUGALL-SHACKLETON, E.A. (2010). Paternal song complexity predicts
1557 offspring sex ratios close to fledging, but not hatching, in song sparrows. *Wilson Journal of*
1558 *Ornithology* **122**, 146-152.

1559

1560 PROKOP, Z.M., MICHALCZYK, L., DROBNIAK, S.M., HERDEGEN, M. & RADWAN, J. (2012). Meta-
1561 analysis suggests choosy females get sexy sons more than “good genes”. *Evolution* **66**, 2665-2673.

1562

1563 PROKUDA, A. & ROFF, D. (2014). The quantitative genetics of sexually selected traits, preferred traits
1564 and preference: a meta-analysis. *Journal of Evolutionary Biology* **27**, 2283-2296.

1565

1566 PRYKE, S.R. & GRIFFITH, S.C. (2009). Genetic incompatibility drives sex allocation and maternal
1567 investment in a polymorphic finch. *Science* **323**, 1605-1607.

1568

1569 PRYKE, S.R. & ROLLINS, L.A. (2012). Mothers adjust offspring sex to match the quality of the rearing
1570 environment. *Proceedings of the Royal Society of London Series B* **279**, 4051-4057.

1571

1572 PRYKE, S.R., ROLLINS, L.A., BUTTEMER, W.A. & GRIFFITH, S.C. (2011). Maternal stress to partner
1573 quality is linked to adaptive offspring sex ratio adjustment. *Behavioral Ecology* **22**, 717-722.

1574

1575 *QUESTIAU, S., ESCARAVAGE, N., EYBERT, M.-C. & TABERLET, P. (2000). Nestling sex ratios in a
1576 population of bluethroats *Luscinia svecica* inferred from AFLP (TM) analysis. *Journal of Avian*
1577 *Biology* **31**, 8-14.

1578

1579 RADFORD, A.N. & BLAKEY, J.K. (2000a). Intensity of nest defence is related to offspring sex ratio in
1580 the great tit *Parus major*. *Proceedings of the Royal Society of London Series B* **267**, 535-538.

1581

1582 *RADFORD, A.N. & BLAKEY, J.K. (2000b). Is variation in brood sex ratios adaptive in the great tit
1583 (*Parus major*)? *Behavioral Ecology* **11**, 294-298.

1584

1585 *RAMSAY, S.M., MENNILL, D.J., OTTER, K.A., RATCLIFFE, L.M. & BOAG, P.T. (2003). Sex allocation
1586 in black-capped chickadees *Poecile atricapilla*. *Journal of Avian Biology* **34**, 134-139.

1587

1588 *RATHBURN, M.K. & MONTGOMERIE R. (2005). Offspring sex ratios correlate with pair-male
1589 condition in a cooperatively breeding fairy-wren. *Behavioral Ecology* **16**, 41-47.

1590

1591 RICE, W.R. (2013). Nothing in genetics makes sense except in light of genomic conflict. *Annual*
1592 *Review of Ecology, Evolution and Systematics* **44**, 217-237.

1593

1594 RICE, W.R. & CHIPPINDALE, A.K. (2001). Intersexual ontogenetic conflict. *Journal of Evolutionary*
1595 *Biology* **14**, 685-693.

1596

1597 RØED, K.H., HOLAND, Ø., MYSTERUD, A., TVERDAL, A., KUMPULA, J. & NIEMINEN, M. (2007). Male
1598 phenotypic quality influences offspring sex ratio in a polygynous ungulate. *Proceedings of the Royal*
1599 *Society of London Series B* **274**, 727-733.

1600

1601 ROMANO, A., AMBROSINI, R., CAPRIOLI, M., BONISOLI-ALQUATI, A. & SAINO, N. (2012). Secondary
1602 sex ratio covaries with demographic trends and ecological conditions in the barn swallow.
1603 *Evolutionary Ecology* **26**, 1041-1053.

1604

1605 ROSENTHAL, R. (1979). The file drawer problem and tolerance for null results. *Psychological Bulletin*
1606 **86**, 638-641.

1607

1608 *ROSIVALL, B., TÖRÖK, J., HASSELQUIST, D. & BENSCH, S. (2004). Brood sex ratio adjustment in
1609 collared flycatchers (*Ficedula albicollis*): results differ between populations. *Behavioral Ecology and*
1610 *Sociobiology* **56**, 346-351.

1611

1612 *ROULIN, A., ALTWEGG, R., JENSEN, H., STEINSLAND, I. & SCHAUB, M. (2010). Sex-dependent
1613 selection on an autosomal melanic female ornament promotes the evolution of sex ratio bias. *Ecology*
1614 *Letters* **13**, 616-626.

1615

1616 RUTKOWSKA, J., DUBIEC, A. & NAKAGAWA, S. (2014). All eggs are made equal: meta-analysis of egg
1617 sexual size dimorphism in birds. *Journal of Evolutionary Biology* **27**, 153-160.

1618

1619 *RUTSTEIN, A.N., GILBERT, L., SLATER, P.J.B. & GRAVES, J.A. (2004). Mate attractiveness and
1620 primary resource allocation in the zebra finch. *Animal Behaviour* **68**, 1087-1094.

1621

1622 *RUTSTEIN, A.N., GORMAN, H.E., ARNOLD, K.E., GILBERT, L., ORR, K.J., ADAM, A., NAGER, R. &
1623 GRAVES, J.A. (2005). Sex allocation in response to paternal attractiveness in the zebra finch.
1624 *Behavioral Ecology* **16**, 763-769.

1625

1626 RUTZ, C. (2012). Brood sex ratio varies with diet composition in a generalist raptor. *Biological*
1627 *Journal of the Linnaean Society* **105**, 937-951.

1628

1629 SAETHER, B.-E., SOLBERG, E.J., HEIM, M., STACY, J.E., JAKOBSEN K.S. & OLSTAD R. (2004).
1630 Offspring sex ratio in moose *Alces alces* in relation to paternal age: an experiment. *Wildlife Biology*
1631 **10**, 51-57.

1632

1633 *SAINO, N., AMBROSINI, R., MARTINELLI, R., CALZA, S., MØLLER, A.P. & PILASTRO, A. (2002).
1634 Offspring sexual dimorphism and sex-allocation in relation to parental age and paternal
1635 ornamentation in the barn swallow. *Molecular Ecology* **11**, 1533-1544.

1636

1637 *SAINO, N., ELLEGREN, H. & MØLLER, A.P. (1999). No evidence for adjustment of sex allocation in
1638 relation to paternal ornamentation and paternity in barn swallows. *Molecular Ecology* **8**, 399-406.

1639

1640 SAINO, N., MARTINELLI, R. & ROMANO, M. (2008). Ecological and phenological covariates of
1641 offspring sex ratio in barn swallows. *Evolutionary Ecology* **22**, 659-674.

1642

1643 *SAINO, N., ROMANO, M., FERRARI, R.P., MARTINELLI, R. & MØLLER, A.P. (2003). Maternal
1644 antibodies but not carotenoids in barn swallow eggs covary with embryo sex. *Journal of Evolutionary*
1645 *Biology* **16**, 516-522.

1646

1647 SARDELL, R.J. & DUVAL, E.H. (2014). Differential allocation in a lekking bird: females lay larger
1648 eggs and are more likely to have male chicks when they mate with less related males. *Proceedings of*
1649 *the Royal Society of London Series B* **281**, 20132386.

1650

1651 *SATO, A. & KARINO, K. (2010). Female control of offspring sex ratios based on male attractiveness
1652 in the guppy. *Ethology* **116**, 524-534.
1653

1654 SATO, A. & KARINO, K. (2011). Offspring sex ratio produced by female guppies in the wild correlates
1655 with sexual ornaments of their sons. *Journal of Ethology* **29**, 221-226.
1656

1657 SCHINO, G. (2004). Birth sex ratio and social rank: consistency and variability between primate
1658 groups. *Behavioral Ecology* **15**, 850-856.
1659

1660 *SCHWARZOVA, L., SIMEK, J., COPPACK, T. & TRYJANOWSKI, P. (2008). Male-biased sex of extra pair
1661 young in the socially monogamous Red-backed Shrike *Lanius collurio*. *Acta Ornithologica* **43**, 235-
1662 240.
1663

1664 SHELDON, B.C. (1998). Recent studies of avian sex ratios. *Heredity* **80**, 397-402.
1665

1666 SHELDON, B.C. (2000). Differential allocation: tests, mechanisms and implications. *Trends in*
1667 *Ecology and Evolution* **15**, 397-402.
1668

1669 *†SHELDON, B.C., ANDERSSON, S., GRIFFITH, S.C., ÖRNBORG, J. & SENDECKA, J. (1999). Ultraviolet
1670 colour variation influences blue tit sex ratios. *Nature* **402**, 874-877.
1671

1672 *SHELDON, B.C. & ELLEGREN, H. (1996). Offspring sex and paternity in the collared flycatcher.
1673 *Proceedings of the Royal Society of London Series B* **263**, 1017-1021.
1674

1675 SHELDON, B.C. & WEST, S.A. (2004). Maternal dominance, maternal condition and offspring sex
1676 ratio in ungulate mammals. *The American Naturalist* **163**, 40-54.
1677

1678 SHUKER, D.M. & WEST, S.A. (2004). Information constraints and the precision of adaptation: sex
1679 ratio manipulation in wasps. *Proceedings of the National Academy of Sciences of the United States of*
1680 *America* **101**, 10363-10367.
1681

1682 SHUSTER, S.M. & WADE, M.J. (2003). *Mating Systems and Strategies*. Princeton University Press,
1683 Princeton, NJ.
1684

1685 SILK, J.B. & BROWN, G.R. (2008). Local resource competition and local resource enhancement shape
1686 primate birth sex ratios. *Proceedings of the Royal Society of London Series B* **275**, 1761-1765.
1687

1688 SILK, J.B., WILLOUGHBY, E. & BROWN, J.R. (2005). Maternal rank and local resource competition do
1689 not predict birth sex ratios in wild baboons. *Proceedings of the Royal Society of London Series B* **272**,
1690 859-864.
1691

1692 SIMMONS, J.P., NELSON, L.D. & SIMONSOHN, U. (2011). False-positive psychology: Undisclosed
1693 flexibility in data collection and analysis allows presenting anything as significant. *Psychological*
1694 *Science* **22**, 1359-1366.
1695

1696 SIMONSOHN, U., NELSON, L.D. & SIMMONS, J.P. (2014). P-curve: a key to the file drawer. *Journal of*
1697 *Experimental Psychology* **143**, 534-547.
1698

1699 *SOMA, M. & OKANOYA, K. (2013). Differential allocation in relation to mate song quality in the
1700 Bengalese finch. *Behaviour* **150**, 1491-1508.
1701

1702 *STAUSS, M., SEGELBACHER, G., TOMIUK, J. & BACHMANN, L. (2005). Sex ratio of *Parus major* and
1703 *P. caeruleus* broods depends on parental condition and habitat quality. *Oikos* **109**, 367-373.
1704

1705 STERNE, J.A.C. & EGGER, M. (2005). Regression method to detect publication and other bias in meta-
1706 analysis. In: *Publication bias in meta-analysis: prevention, assessment and adjustments* (eds. H.
1707 ROTHSTEIN, A.J. SUTTON & M. BORENSTEIN), pp. 99-125. John Wiley & Sons, Chichester, UK.
1708

1709 STUBBLEFIELD, J.W. & ORZACK, S.H. (2013). Resource transfers and evolution: helpful offspring and
1710 sex allocation. *Theoretical Population Biology* **83**, 64-81.
1711

1712 *SVENSSON, E. & NILSSON, J.-A. (1996). Mate quality affects offspring sex ratio in blue tits.
1713 *Proceedings of the Royal Society of London Series B* **263**, 357-361.
1714

1715 *SZÉKELY, T., CUTHILL, I.C., YEZERINAC, S., GRIFFITHS, R. & KIS, J. (2004). Brood sex ratio in the
1716 Kentish plover. *Behavioral Ecology* **15**, 58-62.
1717

1718 *TAFF, C.C., FREEMAN-GALLANT, C.R., DUNN, P.O. & WHITTINGHAM, L.A. (2011). Relationship
 1719 between brood sex ratio and male ornaments depends on male age in a warbler. *Animal Behaviour*
 1720 **81**, 619-625.

1721

1722 TAYLOR, P.D. (1981). Intra-sex and inter-sex sibling interactions as sex-ratio determinants. *Nature*
 1723 **291**, 64-66.

1724

1725 TÖLÜ, C., SAVAS, T., PALA, A. & THOMSEN, H. (2007). Effects of goat social rank on kid gender.
 1726 *Czech Journal of Animal Sciences* **52**, 77-82.

1727

1728 TRIVERS, R.L. & WILLARD, D.E. (1973). Natural selection of parental ability to vary the sex ratio of
 1729 offspring. *Science* **179**, 90-92.

1730

1731 TRNKA, A., PROKOP, P., KASOVA, M., SOBEKOVA, K. & KOCIAN, L. (2012). Hatchling sex ratio and
 1732 female mating status in the great reed warbler, *Acrocephalus arundinaceus* (Aves, Passeriformes):
 1733 further evidence for offspring sex ratio manipulation. *Italian Journal of Zoology* **79**, 212-217.

1734

1735 UNGERFELD, R., GONZÁLEZ-SIERRA, U.T. & PIAGGIO, J. (2008). Reproduction in a semi-captive herd
 1736 of pampas deer *Ozotocerus bezoarticus*. *Wildlife Biology* **14**, 350-357.

1737

1738 VEEN, T., TRÄFF, J., WEISSING, F.J. & SHELDON, B.C. (2009). Reduced costs of mixed-species
 1739 pairings in flycatchers: by-product or female strategy? *Behavioral Ecology and Sociobiology* **63**, 329-
 1740 337.

1741

1742 *VEIGA, J.P., POLO, V. & CORDERO, P.J. (2008). Age dependent sex allocation in the polygynous
 1743 spotless starling. *Evolutionary Ecology* **22**, 167-176.

1744

1745 VIECHTBAUER, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of*
 1746 *Statistical Software* **36**, 1-48.

1747

1748 WALSH B, & BLOWS, M.W. (2009). Abundant genetic variation plus strong selection = multivariate
 1749 genetic constraints: a geometric view of adaptation. *Annual Review of Ecology, Evolution and*
 1750 *Systematics* **40**, 41-59.

1751

1752 WARNER, D.A., LOVERN, M.B. & SHINE, R. (2008). Maternal influences on offspring phenotypes and
 1753 sex ratios in a multi-clutching lizard with environmental sex determination. *Biological Journal of the*
 1754 *Linnean Society* **95**, 256-266.

1755

1756 *WATT, P.J., SHOHEIT, A.J. & RENSHAW, K. (2001). Female choice for good genes and sex-biased
 1757 broods in guppies. *Journal of Fish Biology* **59**, 843-850.

1758

1759 WEATHERHEAD, P.J. (1983). Secondary sex-ratio adjustment in red-winged blackbirds (*Agelaius*
 1760 *phoeniceus*). *Behavioral Ecology and Sociobiology* **12**, 57-61.

1761

1762 WEATHERHEAD, P.J. & ROBERTSON, R.J. (1979). Offspring quality and the polygyny threshold – sexy
 1763 son hypothesis. *The American Naturalist* **113**, 201-208.

1764

1765 WERREN, J.H. (1983). Sex-ratio evolution under local mate competition in a parasitic wasp. *Evolution*
 1766 **37**, 116-124.

1767

1768 WEST, S.A. (2009). *Sex Allocation*. Princeton University Press, Princeton, NJ.

1769

1770 WEST, S.A., HERRE, E.A. & SHELDON, B.C. (2000). The benefits of allocating sex. *Science* **290**, 288-
 1771 290.

1772

1773 WEST, S.A., REECE, S.E. & SHELDON, B.C. (2002). Sex ratios. *Heredity* **88**, 117-124.

1774

1775 WEST, S.A., SHUKER, D.M. & SHELDON, B.C. (2005). Sex-ratio adjustment when relatives interact: a
 1776 test of constraints on adaptation. *Evolution* **59**, 1211-1228.

1777

1778 †WEST, S.A. & SHELDON, B.C. (2002). Constraints in the evolution of sex ratio adjustment. *Science*
 1779 **295**, 1685-1688.

1780

1781 *WESTERDAHL, H., BENSCH, S., HANSSON, B., HASSELQUIST, D. & VON SCHANTZ, T. (1997). Sex ratio
 1782 variation among broods of great reed warblers *Acrocephalus arundinaceus*. *Molecular Ecology* **6**,
 1783 543-548.

1784

1785 WESTNEAT, D.F. & STEWART, I.R.K. (2003). Extra-pair paternity in birds: causes, correlates, and
 1786 conflict. *Annual Review of Ecology, Evolution and Systematics* **34**, 365-396.

1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821

- *WESTNEAT, D.F., STEWART, I.R.K., WOESTE, E.H., GIPSON, J., ABDULKADIR, L. & POSTON, J.P. (2002). Patterns of sex ratio variation in house sparrows. *Condor* **104**, 598-609.
- WHEELWRIGHT, N.T. & SEABURY, R.E. (2003). Fifty:fifty offspring sex ratios in savannah sparrows (*Passerculus sandwichensis*). *The Auk* **120**, 171-179.
- *WHITTINGHAM, L.A. & DUNN, P.O. (2000). Offspring sex ratios in tree swallows: females in better condition produce more sons. *Molecular Ecology* **9**, 1123-1129.
- *WIEBE, K.L. & BORTOLOTTI, G.R. (1992). Facultative sex-ratio manipulation in American kestrels. *Behavioral Ecology and Sociobiology* **30**, 379-386.
- *WILK, T., CICHON, M. & WOLFF, K. (2008). Lack of evidence for improved immune response of extra-pair nestlings in collared flycatcher *Ficedula albicollis*. *Journal of Avian Biology* **39**, 546-552.
- WOOD, W., KRESSEL, L., JOSHI, P.D. & LOUIE, B. (2014). Meta-analysis of menstrual cycle effects on women's mate preferences. *Emotion Review* **6**, 229-249.
- WYMAN, M.J. & ROWE, L. (2014). Male bias in distributions of additive genetic, residual, and phenotypic variances of shared traits. *The American Naturalist* **184**, 326-337.
- *YAMAGUCHI, N., KAWANO, K.K., EGUCHI, K. & YAHARA, T. (2004). Facultative sex ratio adjustment in response to male tarsus length in the varied tit *Parus varius*. *Ibis* **146**, 108-113.
- *ZANN, R. & RUNCIMAN, D. (2003). Primary sex ratios in zebra finches: no evidence for adaptive manipulation in wild and semi-domesticated populations. *Behavioral Ecology and Sociobiology* **54**, 294-302.
- *ZIELINSKA, M., DUBIEC, A. & ZIELINSKI, P. (2010). Offspring sex ratio skew in the sexually monomorphic house martin *Delichon urbicum*. *Journal of Avian Biology* **41**, 591-596.

1822 **VIII. SUPPORTING INFORMATION**

1823 Additional supporting information may be found in the online version of this article,

1824 DOI: 10.1111/brv.12220

1825

1826 **Fig. S1.** The phylogenetic trees used for the phylogenetic meta-analysis and meta-regression
1827 analysis: (1) a tree with Ericson backbone and (2) a tree with Hackett backbone.

1828

1829 **Fig. S2.** Forest plot of the regression coefficients of the phylogenetic meta-regression with the
1830 Hackett tree.

1831

1832 **Table S1.** Results of phylogenetic (random-effects) meta-analyses using the Hackett tree.

1833

1834 **Table S2.** Results of multilevel and phylogenetic (random-effects) meta-analyses, incorporating
1835 correlations among effect sizes that come from the same set of individuals.

1836

1837 **Table S3.** Summary of the 25 studies excluded on full-text screening.

1838

1839 **Appendix S1.** Literature search and selection procedure.

1840

1841 **Appendix S2.** EndNote library (SexRatioAdjustment.enlp): all unique records obtained from the *Web*
1842 *of Science* literature searches outlined in Appendix S1.

1843

1844 **Appendix S3.** Raw data file (SexRatioAdjustmentData.xlsx): all effect sizes included in the meta-
1845 analyses.


1846

1847 **Figures**

1848

1849 **Fig. 1.** Papers published per year since Burley (1981) on adjustment of offspring sex ratios in
1850 response to paternal traits. Only papers eligible for inclusion in our meta-analyses are shown; four
1851 included papers from 2014 are not shown as our search only encompassed half of this year.


1852



1853

1854


1855 **Fig. 2.** Preferred reporting items for systematic reviews and meta analyses (PRISMA) flow chart of
 1856 our literature search and data collection process (see <http://www.prisma-statement.org/>).
 1857




1858
 1859
 1860

Fig. 3. Forest plots of (A) the meta-analytic means of the eight random-effects meta-analyses, the multilevel meta-analysis, and the phylogenetic meta-analysis (the latter two used the avian-only data set; the phylogenetic analysis used the Ericson backbone tree), (B) the regression coefficients of the multilevel meta-regression, and (C) of the phylogenetic meta-regression. EPP, extra-pair paternity; k , number of effect sizes.

1866



1867 **Fig. 4.** Funnel plots (effect sizes plotted against their corresponding precision, the inverse of the
1868 standard error) of (A–H) the eight random-effects meta-analyses, (I) the multilevel meta-analysis and
1869 (J) the phylogenetic multilevel meta-analysis (Ericson backbone tree). Plots for the multilevel
1870 analyses (I, J) use the meta-analytic residuals (see Nakagawa & Santos, 2012); these analyses used
1871 the avian-only data set. EPP, extra-pair paternity.



1873
1874

Table 1. Summary of 93 papers coded for the meta-analyses. *N* studies reports the number of independent populations/experiments for which the relationship between brood sex ratios and some measure of male attractiveness was tested. *N* broods gives the range of sample sizes for each study; even where a paper contained only one study, different numbers of broods may have contributed to the tests for different measured traits. For EPP (extra-pair paternity): ‘within broods’ compares the sex ratio of offspring sired by a female’s social and extra-pair mate, within the same brood; ‘between broods’ compares the sex ratio of broods with and without any extra-pair offspring; ‘across all broods’ compares the probability of being male for individual offspring sired by a within- or extra-pair male.

Reference	Species	<i>N</i> studies	Measures of attractiveness tested	<i>N</i> broods
Abroe <i>et al.</i> (2007)	Common yellowthroat, <i>Geothlypis trichas</i>	1	EPP: within broods; Ornament: mask size, bib size, bib colour; Size	69–138
Addison <i>et al.</i> (2008)	Rhinoceros auklet, <i>Cerorhinca monocerata</i>	1	Size	25
Aparicio & Cordero (2001)	Lesser kestrel, <i>Falco naumanni</i>	1	Condition; Ornament: wing patch size	75
Bell <i>et al.</i> (2014)	Blue tit, <i>Cyanistes caeruleus</i>	2	Condition	37, 41
Benito <i>et al.</i> (2013)	Common tern, <i>Sterna hirundo</i>	1	Behaviour: arrival date; Size	81
Blanchard <i>et al.</i> (2007)	Wandering albatross, <i>Diomedea exulans</i>	1	Condition; Size	63
Bowers <i>et al.</i> (2013)	Collared flycatcher, <i>Ficedula albicollis</i>	1	Age; EPP: across all broods; Ornament: wing patch size, forehead patch size	160
Bradbury <i>et al.</i> (1997)	European starling, <i>Sturnus vulgaris</i>	1	Age; Size	108
Burley (1981)	Zebra finch, <i>Taeniopygia guttata</i>	1	Ornament: leg band colour	20
Burley (1986)	Zebra finch, <i>Taeniopygia guttata</i>	1	Ornament: leg band colour	18
Calsbeek & Sinervo (2004)	Side-blotched lizard, <i>Uta stansburiana</i>	2	Size	33, 41
Cockburn & Double (2008)	Superb fairy wren, <i>Malurus cyaneus</i>	1	Condition; EPP: across all broods	993
Correa <i>et al.</i> (2011)	Japanese quail, <i>Coturnix japonica</i>	1	Behaviour: strutting, neck holding, mounting	26
Cotton <i>et al.</i> (2014)	Stalk-eyed fly, <i>Teleopsis dalmanni</i>	1	Ornament: eyespan, relative eyespan; Size	130
Cox <i>et al.</i> (2011)	Brown anole, <i>Anolis sagrei</i>	1	Condition	67
Cunningham & Russell (2001)	Mallard, <i>Anas platyrhynchos</i>	1	Female preference	32
Debiak <i>et al.</i> (2014)	Common loon, <i>Gavius immer</i>	1	Size	14
Delhey <i>et al.</i> (2007)	Blue tit, <i>Cyanistes caeruleus</i>	4	Ornament: crown brightness, chroma, hue, UV chroma; Size	20, 22, 40, 70
Delmore <i>et al.</i> (2008)	Tree swallow, <i>Tachycineta bicolor</i>	1	Age; Condition; EPP: between broods; Size	50
Dietrich-Bischoff <i>et al.</i> (2006)	Coal tit, <i>Parus ater</i>	1	Age; EPP: within broods, between broods; Ornament: bib saturation, bib size; Size	235–457
Dolan <i>et al.</i> (2009)	Eastern kingbird, <i>Tyrannus tyrannus</i>	1	Behaviour: song latency, song rate; EPP: between broods; Size	42–72
Dowling & Mulder (2006)	Red-capped robin, <i>Petroica goodenovii</i>	1	EPP: within broods; Ornament: breast & forehead patch brightness, hue, chroma; Size	44
Dreiss <i>et al.</i> (2006)	Blue tit, <i>Cyanistes caeruleus</i>	1	Behaviour: song characteristics; EPP: within broods; Ornament: crown brightness, hue, UV chroma; Size	84
Drullion & Dubois (2011)	Zebra finch, <i>Taeniopygia guttata</i>	1	Female preference	6

Reference	Species	N studies	Measures of attractiveness tested	N broods
Du & Lu (2010)	Tibetan ground tit, <i>Pseudopodoces humilis</i>	2	EPP: between broods	19, 58
Ellegren <i>et al.</i> (1996)	Collared flycatcher, <i>Ficedula albicollis</i>	1	Ornament: forehead patch size; Size	79
Fargallo <i>et al.</i> (2004)	Chinstrap penguin, <i>Pygoscelis antarctica</i>	1	Behaviour: nest size; Size	37–53
Ferree (2007)	Dark-eyed junco, <i>Junco hyemalis</i>	1	Ornament: tail patch size; Size	32
Green (2002)	Brown thornbill, <i>Acanthiza pusilla</i>	1	Size	104
Griffith <i>et al.</i> (2003)	Blue tit, <i>Cyanistes caeruleus</i>	3	Age; Ornament: plumage brightness, UV chroma	15–16, 22–23, 50–57
Grindstaff <i>et al.</i> (2001)	Dark-eyed junco, <i>Junco hyemalis</i>	1	Female preference	48
Helle <i>et al.</i> (2008)	Field vole, <i>Microtus agrestis</i>	1	Condition	28
Hjernquist <i>et al.</i> (2009)	Collared flycatcher, <i>Ficedula albicollis</i>	1	Condition; Ornament: forehead patch size	74
House <i>et al.</i> (2011)	Dung beetle, <i>Onthophagus taurus</i>	1	Size	50
Husby <i>et al.</i> (2006)	House sparrow, <i>Passer domesticus</i>	1	Age; Ornament: badge size; Size	15
Johnson <i>et al.</i> (2009)	House wren, <i>Troglodytes aedon</i>	1	EPP: within broods, between broods	476
Karino <i>et al.</i> (2006)	Guppy, <i>Poecilia reticulata</i>	1	Ornament: tail length	74
Karino & Sato (2009)	Guppy, <i>Poecilia reticulata</i>	1	Ornament: orange spot brightness; Female preference	37
Katsuki <i>et al.</i> (2012)	Broad-horned flour beetle, <i>Gnathocerus cornutus</i>	1	Ornament: mandible length	50
Kempenaers <i>et al.</i> (1997)	Blue tit, <i>Cyanistes caeruleus</i>	1	EPP: within broods, between broods	57–141
Kölliker <i>et al.</i> (1999)	Great tit, <i>Parus major</i>	1	Age; Ornament: breast stripe size; Size	57
Korpimäki <i>et al.</i> (2000)	Kestrel, <i>Falco tinnunculus</i>	1	Condition; Size	80
Korsten <i>et al.</i> (2006)	Blue tit, <i>Cyanistes caeruleus</i>	2	Ornament: UV chroma	26, 44
Kraaijeveld <i>et al.</i> (2007)	Black swan, <i>Cygnus atratus</i>	1	EPP: across all broods; Ornament: number of curled wing feathers; Size	36
Lee & Hatchwell (2011)	Vinous-throated parrotbill, <i>Paradoxomis webbianus</i>	1	EPP: between broods; Size	36
Leech <i>et al.</i> (2001)	Blue tit, <i>Cyanistes caeruleus</i>	1	Age; Condition; EPP: within broods; Size	103–154
Leitner <i>et al.</i> (2006)	Canary, <i>Serinus canaria</i>	1	Behaviour: song components	18
Leonard & Weatherhead (1996)	Red junglefowl (chicken), <i>Gallus gallus</i>	3	Behaviour: social dominance rank	31, 116, 37
Lessells <i>et al.</i> (1996)	Great tit, <i>Parus major</i>	1	Age; Size	52–92
Limboung <i>et al.</i> (2013)	Blue tit, <i>Cyanistes caeruleus</i>	1	Age; Ornament: UV chroma	33
Long & Pischedda (2005)	Fruit fly, <i>Drosophila melanogaster</i>	2	Age	35–131, 384
Lu <i>et al.</i> (2013)	White-rumped snowfinch, <i>Montifringilla taczanowskii</i>	2	Size	21
Magrath <i>et al.</i> (2002)	Fairy martin, <i>Petrochelidon ariel</i>	1	Age; Condition; EPP: between broods; Size	29–56
Neto <i>et al.</i> (2011)	Savi's warbler, <i>Locustella luscinioides</i>	1	EPP: between broods; Size	102
Nishiumi (1998)	Great reed warbler, <i>Acrocephalus arundinaceus</i>	1	Behaviour: arrival date	50
Oddie & Reim (2002)	Great tit, <i>Parus major</i>	1	Age; Condition; Size	143–173
Olsson <i>et al.</i> (2005b)	Sand lizard, <i>Lacerta agilis</i>	1	Ornament: badge size	58
Pariser <i>et al.</i> (2012)	Zebra finch, <i>Taeniopygia guttata</i>	1	Ornament: leg band colour	21
Parker (2005)	Red junglefowl (chicken), <i>Gallus gallus</i>	1	Ornament: comb size	76

Reference	Species	N studies	Measures of attractiveness tested	N broods
Perlut <i>et al.</i> (2014)	Savannah sparrow, <i>Passerculus sandwichensis</i>	1	EPP: between broods; Size	26–48
Pike & Petrie (2005)	Peafowl, <i>Pavo cristatus</i>	1	Ornament: number of eyespot feathers in tail	21
Polo <i>et al.</i> (2004)	Spotless starling, <i>Sturnus unicolor</i>	1	Behaviour: nest decoration	149
Potvin & MacDougall-Shackleton (2010)	Song sparrow, <i>Melospiza melodia</i>	1	Behaviour: song repertoire	55
Questiau <i>et al.</i> (2000)	Bluethroat, <i>Luscinia svecica</i>	1	Age; Ornament: throat patch size; Size	27–39
Radford & Blakey (2000b)	Great tit, <i>Parus major</i>	1	Age; Condition; Ornament: breast stripe size; Size	84
Ramsay <i>et al.</i> (2003)	Black-capped chickadee, <i>Poecile atricapilla</i>	1	Behaviour: social dominance rank; EPP: between broods	150–174
Rathburn & Montgomerie (2005)	White-winged fairy wren, <i>Malurus leucopterus</i>	1	Condition; Ornament: nuptial plumage	79
Rosivall <i>et al.</i> (2004)	Collared flycatcher, <i>Ficedula albicollis</i>	1	Age; Ornament: forehead patch size, wing patch size; Size	54–57
Roulin <i>et al.</i> (2010)	Barn owl, <i>Tyto alba</i>	1	Ornament: spot size	44
Rutstein <i>et al.</i> (2004)	Zebra finch, <i>Taeniopygia guttata</i>	1	Ornament: leg band colour	32
Rutstein <i>et al.</i> (2005)	Zebra finch, <i>Taeniopygia guttata</i>	1	Ornament: leg band colour	101
Saino <i>et al.</i> (1999)	Barn swallow, <i>Hirundo rustica</i>	1	Condition; EPP: across all broods; Ornament: tail length; Size	19–54
Saino <i>et al.</i> (2002)	Barn swallow, <i>Hirundo rustica</i>	1	Age; Ornament: tail length	76
Saino <i>et al.</i> (2003)	Barn swallow, <i>Hirundo rustica</i>	1	Ornament: tail length	46
Sato & Karino (2010)	Guppy, <i>Poecilia reticulata</i>	1	Female preference	30
Schwarzova <i>et al.</i> (2008)	Red-backed shrike, <i>Lanius collurio</i>	1	EPP: across all broods	15
Sheldon & Ellegren (1996)	Collared flycatcher, <i>Ficedula albicollis</i>	1	EPP: within broods	19
Sheldon <i>et al.</i> (1999)	Blue tit, <i>Cyanistes caeruleus</i>	2	Age; Ornament: plumage hue, brightness, UV chroma	20, 57
Soma & Okanoya (2013)	Bengalese finch, <i>Lonchura striata</i>	1	Behaviour: repertoire, song duration, song linearity	29
Stauss <i>et al.</i> (2005)	(a) Great tit, <i>Parus major</i>	2	Condition	(a) 95
	(b) Blue tit, <i>Cyanistes caeruleus</i>			(b) 88
Svensson & Nilsson (1996)	Blue tit, <i>Cyanistes caeruleus</i>	1	Age	33
Szekely <i>et al.</i> (2004)	Kentish plover, <i>Charadrius alexandrinus</i>	1	Age; Size	59
Taff <i>et al.</i> (2011)	Common yellowthroat, <i>Geothlypis trichas</i>	2	Age; EPP: within broods; Ornament: plumage yellow & UV brightness, carotenoid chroma, mask size, bib size; Size	27–71, 44
Veiga <i>et al.</i> (2008)	Spotless starling, <i>Sturnus unicolor</i>	3	Behaviour: nest decoration	34, 40, 135
Watt <i>et al.</i> (2001)	Guppy, <i>Poecilia reticulata</i>	1	Ornament: orange spot size; Female preference; Size	17–24
Westerdahl <i>et al.</i> (1997)	Great reed warbler, <i>Acrocephalus arundinaceus</i>	1	Behaviour: song repertoire; EPP: within broods; Size	9–63
Westneat <i>et al.</i> (2002)	House sparrow, <i>Passer domesticus</i>	1	Age; Condition; Ornament: bib size; Size	93–142
Whittingham & Dunn (2000)	Tree swallow, <i>Tachycineta bicolor</i>	1	Condition	40

Reference	Species	N studies	Measures of attractiveness tested	N broods
Wiebe & Bortolotti (1992)	American kestrel, <i>Falco sparverius</i>	1	Size	87
Wilk <i>et al.</i> (2008)	Collared flycatcher, <i>Ficedula albicollis</i>	1	EPP: across all broods	78
Yamaguchi <i>et al.</i> (2004)	Varied tit, <i>Parus varius</i>	1	Ornament: forehead & breast patch size; Size	59
Zann & Runciman (2003)	Zebra finch, <i>Taeniopygia guttata</i>	3	Condition; Ornament: breast band size, leg band colour; Size	20, 24, 68
Zielinska <i>et al.</i> (2010)	House martin, <i>Delichon urbicum</i>	1	Condition; Ornament: rump patch colour & size; Size	129

Table 2. Results from the random-effects meta-analyses for the eight different trait categories; k = number of effect sizes, m = number of species and n = number of broods in each data set used for these meta-analyses (for the descriptions of Dataset, Q and I^2 , see text Section II); EPP, extra-pair paternity.

Trait	Dataset	k	m	n	Mean (Z_r)	Lower CI (2.5%)	Upper CI (97.5%)	z value	$P(z)$	Q	$d.f. (Q)$	$P(Q)$	I^2 (%)
All	1	81	49	7512	0.096	0.057	0.134	4.815	<0.001	177.40	80	<0.001	56.20
	2	84	50	7807	0.072	0.040	0.104	4.432	<0.001	136.70	83	<0.001	37.07
	3	94	57	8788	0.061	0.033	0.089	4.238	<0.001	138.35	93	0.002	30.02
	4	94	57	8788	0.039	0.012	0.067	2.817	0.005	135.46	93	0.003	26.77
Age	1	9	6	1053	0.015	-0.096	0.127	0.273	0.785	19.60	8	0.012	58.81
Behaviour	1	13	11	1159	0.078	-0.005	0.161	1.849	0.064	23.72	12	0.022	43.21
Condition	1	14	12	1908	0.062	-0.046	0.170	1.118	0.264	38.59	13	<0.001	73.29
EPP	1	16	14	3159	0.016	-0.021	0.052	0.851	0.395	15.26	15	0.433	0.000
Ornament	1	37	18	2394	0.073	0.017	0.130	2.534	0.011	61.86	36	0.005	42.56
	2	38	18	2418	0.072	0.016	0.128	2.528	0.012	61.94	37	0.006	41.31
	3	43	21	2911	0.060	0.012	0.109	2.451	0.014	63.61	42	0.017	34.68
	4	43	21	2911	0.042	-0.006	0.089	1.708	0.088	64.32	42	0.015	33.26
Preference	1	6	4	172	0.148	-0.180	0.477	0.885	0.376	17.26	5	0.004	73.43
Size	1	30	27	1757	0.097	0.033	0.162	2.950	0.003	49.02	29	0.011	42.67
	2	31	28	1865	0.093	0.031	0.155	2.926	0.003	50.10	30	0.012	41.86
	3	44	36	3207	0.060	0.018	0.102	2.778	0.005	57.30	43	0.071	25.75
	4	44	36	3207	0.025	-0.020	0.069	1.092	0.275	61.98	43	0.030	32.41

Table 3. Results of Egger's regression tests and trim-and-fill (TAF) tests; * indicates missing effect sizes in the unexpected direction (for the descriptions of Q and I^2 , see text Section II); EPP, extra-pair paternity.

Trait	t : Egger	$d.f.$ (t)	$P(t)$	Missing k : TAF	Mean (Zr)	Lower CI (2.5%)	Upper CI (97.5%)	z value	$P(z)$	Q	$d.f. (Q)$	$P(Q)$	I^2 (%)
All	3.450	79	0.001	17	0.034	-0.012	0.080	1.443	0.149	287.08	97	<0.001	72.99
Age	3.650	7	0.008	4	-0.102	-0.239	0.036	-1.447	0.148	44.40	12	<0.001	80.17
Behaviour	0.918	11	0.378	0	—	—	—	—	—	—	—	—	—
Condition	0.949	12	0.361	3	-0.009	-0.129	0.111	-0.143	0.886	63.89	16	<0.001	81.71
EPP	0.293	14	0.774	0	—	—	—	—	—	—	—	—	—
Ornament	1.515	35	0.139	7	0.029	-0.031	0.089	0.935	0.350	92.54	43	<0.001	53.75
Preference	-0.873	4	0.432	2*	0.304	-0.046	0.654	1.701	0.089	27.09	7	<0.001	78.40
Size	-0.984	28	0.334	7*	0.151	0.087	0.215	4.618	<0.001	74.02	36	<0.001	52.33

Table 4. Results of multilevel and phylogenetic (random-effects) meta-analyses; k = number of effect sizes, m = number of species and n = number of experiments/populations (or studies) in each data set used for these meta-analyses (for the descriptions of Dataset, H^2 and I^2 , see the text Section II).

Type	Dataset (avian)	k	m	n	Mean (Z_r)	Lower CI (2.5%)	Upper CI (97.5%)	I^2_{study} (%)	I^2_{species} (%)	$I^2_{\text{effect size}}$ (%)	$I^2_{\text{phylogeny}}$ (%)	I^2_{total} (%)	H^2 (%)
Multilevel	1	195	42	85	0.064	0.030	0.103	21.03	6.03	24.50	–	51.56	–
(non-phylogenetic)	2	205	43	88	0.058	0.026	0.093	17.53	5.84	23.79	–	47.16	–
	3	254	48	99	0.041	0.017	0.066	11.04	5.59	12.01	–	28.64	–
	4	254	48	99	0.025	–0.002	0.052	12.37	6.08	18.84	–	37.29	–
Phylogenetic	1	195	42	85	0.074	0.005	0.155	16.26	5.45	23.21	10.88	55.79	9.82
	2	205	43	88	0.069	–0.009	0.142	15.34	5.06	21.40	10.47	52.26	9.77

Table 5. Factors proposed to affect the relative fitness of sons and daughters differentially, leading to adaptive sex ratio adjustment strategies. Examples are given of empirical studies that report a significant effect of the factor; note that for all factors we also located studies that find no relationship. References to meta-analyses are provided where possible.

Influencing factor		References
Maternal condition		Trivers & Willard (1973); Clutton-Brock <i>et al.</i> (1984); Kruuk <i>et al.</i> (1999); meta-analyses: Cameron (2004); Sheldon & West (2004)
Mate attractiveness/quality		Burley (1981, 1986) (and see Table 1, this study); meta-analyses: West & Sheldon (2002); Ewen <i>et al.</i> (2004); Cassey <i>et al.</i> (2006)
Local resource enhancement (cooperative breeders: number of helpers)		Taylor (1981); Gowaty & Lennartz (1985); meta-analyses: West & Sheldon (2002); Griffin <i>et al.</i> (2005); Stubblefield & Orzack (2013)
Local resource competition		Taylor (1981); Cockburn <i>et al.</i> (1985); meta-analysis: Silk & Brown (2008)
Local mate competition		Hamilton (1967); Werren (1983); Shuker & West (2004); meta-analysis: West <i>et al.</i> (2005)
Resource quality/availability:	Diet quality, composition	Warner <i>et al.</i> (2008); Pryke & Rollins (2012); Rutz (2012)
	Habitat/territory quality	Romano <i>et al.</i> (2012); Bell <i>et al.</i> (2014)
	Host size for parasitoid larval development	Godfray (1994); meta-analysis: West & Sheldon (2002)
Seasonal effects:	Time of breeding/laying date	Saino <i>et al.</i> (2008); Graham <i>et al.</i> (2011); Barclay (2012); meta-analysis: Ewen <i>et al.</i> (2004); Cassey <i>et al.</i> (2006)
	Rainfall	Berkeley & Linklater (2010)
Laying/hatch order		Cichon <i>et al.</i> (2003); Lezalova <i>et al.</i> (2005); Lislevand <i>et al.</i> (2005)
Clutch size		Benito <i>et al.</i> (2013); Bowers <i>et al.</i> (2014)
Social dominance rank (maternal)		Nevison (1997); Maestripieri (2002); meta-analyses: Brown & Silk (2002); Schino (2004); Silk <i>et al.</i> (2005)
Parental relatedness/compatibility		Pryke & Griffith (2009); Sardell & DuVal (2014)
Parental age/breeding experience/pair bond duration		Green (2002); Benito <i>et al.</i> (2013)
Female mating status/extent of polygyny (sole/primary/secondary female, harem size)		Weatherhead (1983); Nishiumi (1998); Trnka <i>et al.</i> (2012)